

GRADIENT ESTIMATES FOR THE CONDUCTIVITY PROBLEMS

BY SHITING BAO

A dissertation submitted to the
Graduate School—New Brunswick
Rutgers, The State University of New Jersey
in partial fulfillment of the requirements

for the degree of

Doctor of Philosophy

Graduate Program in Mathematics

Written under the direction of

YanYan Li

and approved by

New Brunswick, New Jersey

May, 2008

ABSTRACT OF THE DISSERTATION

Gradient Estimates for the Conductivity Problems

by SHITING BAO

Dissertation Director: YanYan Li

We establish both upper and lower bounds of the gradient estimates for solutions to the perfect conductivity problem in the case where perfect (stiff) conductors are closely spaced inside an open bounded domain. These results give the optimal blow-up rates of the stress for conductors with arbitrary shape and in all dimensions. While for solutions to the insulated conductivity problem we also obtain an upper bound of the gradient estimates.

Acknowledgements

As a student at Rutgers for six years, I have received a lot of strong support and encouragement from professors, family, and friends. I would like to take this opportunity to express my thanks to them.

First and foremost, I want to give my tremendous thanks to my advisor, Professor YanYan Li who opened the door of science in front of me, for his inspiration, encouragement, invaluable guidance and unfailing patience throughout my Ph.D study. Working with him is my most valuable experience and priceless treasures to my future career. I really and deeply appreciate him.

I would like to thank my thesis committee members Professor Zheng-Chao Han, Professor Hyeonbae Kang and Professor Michael Vogelius for their helps during my writing of this dissertation and comments during my graduate career. My appreciation also goes to Professor Haim Brezis for his interest in our work and his strong support.

Thanks also go to Biao Yin for many discussions on mathematics. I appreciate Professor YanYan Li and Biao Yin allowing me to include our joint work in my thesis.

I want to acknowledge to my fellows and friends Qinian Jin, Huaoyuan Xu, Liming Wang, Ren Guo, Luc Nguyen, Yuan Zhang, Jin Wang, Ming Shi, Nan Li, Yuan Yuan, Tianling Jin, Ke Wang, Yu Wang, Haigang Li, and many other friends at Rutgers, for their helps and encouragement during my six years graduate student life here. Thanks go to my friends outside of Rutgers for their constant support and belief in me.

I owe a great deal to Rutgers University and the Department of Mathematics for their support and assistance during my graduate study.

Finally, I would like to express my appreciation to my family, for supporting me everyday in these years and special thanks to my husband, Ming Chen, for his friendship, his love, his trust, and much much more. I am forever thankful.

Dedication

To my grandparents QiRong Bao, YunMei Ling, DuoYun Zhang, and QuanYun Li.

To my parents YingQi Bao, and WeiZhen Zhang.

To my husband Ming Chen.

Table of Contents

Abstract	ii
Acknowledgements	iii
Dedication	iv
1. Introduction	1
2. The perfect conductivity problem with two inclusions	11
2.1. Proof of Theorem 1.1 and 1.2	11
2.1.1. Estimate of $ C_1 - C_2 $	15
2.1.2. Estimate of $ a_{11} - \alpha a_{12} $	17
2.1.3. Estimate of $ b_1 - \alpha b_2 $	21
2.2. Estimate of $ Q_\varepsilon[\varphi] $	23
2.3. Proof of Theorem 1.3 and 1.4	30
2.4. Appendix A: Some elementary results for the perfect conductivity problem	32
3. The perfect and insulated conductivity problems with multiple inclusions	39
3.1. The perfect conductivity problem with multiple inclusions	39
3.1.1. A Matrix Theorem	40
3.1.2. Proof of Theorem 1.5	50
3.2. The insulated conductivity problem	55
3.3. Appendix B: Some elementary results for the insulated conductivity problem	62
References	69
Vita	72

Chapter 1

Introduction

The thesis consists of two parts. The first part concerns the gradient estimates for solutions to the perfect conductivity problem in the case where two perfect (stiff) conductors are closely spaced inside an open bounded domain. The second part studies the gradient estimates for solutions to both the perfect and insulated conductivity problems where there are multiple inclusions. The results in the first part, Chapter 2, have appeared as a preprint in [11]; and the results in the second part, Chapter 3, have appeared as a preprint in [12].

Let Ω be a bounded open set in \mathbb{R}^n with $C^{2,\alpha}$ boundary, $n \geq 2$, $0 < \alpha < 1$, D_1 and D_2 be two bounded strictly convex open subsets in Ω with $C^{2,\alpha}$ boundaries which are ε apart and far away from $\partial\Omega$, i.e.

$$\begin{aligned} \overline{D_1}, \overline{D_2} \subset \Omega, \quad \text{the principal curvature of } \partial D_1, \partial D_2 \geq \kappa_0, \\ \varepsilon := \text{dist}(D_1, D_2) > 0, \quad \text{dist}(D_1 \cup D_2, \partial\Omega) > r_0, \quad \text{diam}(\Omega) < \frac{1}{r_0}, \end{aligned} \tag{1.1}$$

where $\kappa_0, r_0 > 0$ are universal constants independent of ε . We will assume that the $C^{2,\alpha}$ norms of ∂D_i are bounded by some constant independent of ε . This implies $\text{diam}(D_i) \geq r_0^*$ for some universal constant $r_0^* > 0$ independent of ε . We denote

$$\tilde{\Omega} := \Omega \setminus \overline{D_1 \cup D_2}.$$

Given $\varphi \in C^2(\partial\Omega)$, consider the following scalar equation with Dirichlet boundary condition:

$$\begin{cases} \text{div}(a_k(x) \nabla u_k) = 0 & \text{in } \Omega, \\ u_k = \varphi & \text{on } \partial\Omega, \end{cases} \tag{1.2}$$

where

$$a_k(x) = \begin{cases} k \in (0, \infty) & \text{in } D_1 \cup D_2, \\ 1 & \text{on } \Omega \setminus \overline{D_1 \cup D_2}. \end{cases} \tag{1.3}$$

It is well known that there exists a unique solution $u_k \in H^1(\Omega)$ of the above equation, which is also the minimizer of I_k on $H_\varphi^1(\Omega)$, where

$$H_\varphi^1(\Omega) := \{u \in H^1(\Omega) \mid u = \varphi \text{ on } \partial\Omega\}, \quad I_k[v] := \frac{1}{2} \int_{\Omega} a_k |\nabla v|^2.$$

As explained in the introduction of [26], the above equation in dimension $n = 2$ can be used as a simple model in the study of composite media with closely spaced interfacial boundaries. For this purpose, the domain Ω would model the cross-section of a fiber-reinforced composite, D_1 and D_2 would represent the cross-sections of the fibers, $\tilde{\Omega}$ would represent the matrix surrounding the fibers, and the shear modulus of the fibers would be k and that of the matrix would be 1. Equation (1.2) is then obtained by using a standard model of anti-plane shear, and the solution u_k represents the out of plane elastic displacement. The most important quantities from an engineering point of view are the stresses, in this case represented by ∇u_k .

It is well known that the solution u_k satisfies $\|u_k\|_{C^{2,\alpha}(D_i)} < \infty$. In fact, if ∂D_1 and ∂D_2 are $C^{m,\alpha}$, we have $\|u_k\|_{C^{m,\alpha}(D_i)} < \infty$. Such results do not require D_i to be convex and hold for general elliptic systems with piecewise smooth coefficients; see e.g. theorem 9.1 in [26] and proposition 1.6 in [25]. For a fixed $0 < k < \infty$, the $C^{m,\alpha}(D_i)$ -norm of the solution might tend to infinity as $\varepsilon \rightarrow 0$. Babuska, Anderson, Smith and Levin [9] were interested in linear elliptic systems of elasticity arising from the study of composite material. They observed numerically that, for solution u to certain homogeneous isotropic linear systems of elasticity, $\|\nabla u\|_{L^\infty}$ is bounded independently of the distance ε between D_1 and D_1 . Bonnetier and Vogelius [14] proved this in dimension $n = 2$ for the solution u_k of (1.2) when D_1 and D_2 are two unit balls touching at a point. This result was extended by Li and Vogelius in [26] to general second order elliptic equations with piecewise smooth coefficients, where stronger $C^{1,\beta}$ estimates were established. The $C^{1,\beta}$ estimates were further extended by Li and Nirenberg in [25] to general second order elliptic systems including systems of elasticity. For higher derivative estimates, e.g. an ε -independent L^∞ -estimate of second derivatives of u_k in D_1 , we draw attention of readers to the open problem on page 894 of [25]. In [26] and [25], the ellipticity constants are assumed to be away from 0 and ∞ . If we allow ellipticity constants to deteriorate, the situation is different. It has been shown in various papers, see

e.g. [15] and [27], that when $k = \infty$ the L^∞ -norm of ∇u_k for the solution u_k of equation (1.2) generally becomes unbounded as ε tends to zero. The rate at which the L^∞ norm of the gradient of a special solution has been shown in [15] to be $\varepsilon^{-1/2}$. There have been many papers, preprints, expository articles, and works in preparation, on the subject and related ones, see, e.g., [1], [2], [5], [8], [10], [16], [19], [21], [22], [23], [24], [29], [30], [31], [3], [4], [17], [18], [20], [28], [13] and [32].

Part I. In this part, we consider the perfect conductivity problem, where $k = +\infty$.

The perfect conductivity problem is described as follows:

$$\left\{ \begin{array}{ll} \Delta u = 0 & \text{in } \tilde{\Omega}, \\ u|_+ = u|_- & \text{on } \partial D_1 \cup \partial D_2, \\ \nabla u \equiv 0 & \text{in } D_1 \cup D_2, \\ \int_{\partial D_i} \frac{\partial u}{\partial \nu} \Big|_+ = 0 & (i = 1, 2), \\ u = \varphi & \text{on } \partial \Omega. \end{array} \right. \quad (1.4)$$

where

$$\frac{\partial u}{\partial \nu} \Big|_+ := \lim_{t \rightarrow 0^+} \frac{u(x + t\nu) - u(x)}{t}.$$

Here and throughout this thesis ν is the outward unit normal to the domain and the subscript \pm indicates the limit from outside and inside the domain, respectively.

The existence and uniqueness of solutions to equation (1.4) are well known, see the Appendix. Moreover, the solution $u \in H^1(\Omega)$ is the weak limit of the solutions u_k to equations (1.2) as $k \rightarrow +\infty$. It can be also described as the unique function which has the “least energy” in appropriate functional space, defined as $I_\infty[u] = \min_{v \in \mathcal{A}} I_\infty[v]$, where

$$I_\infty[v] := \frac{1}{2} \int_{\tilde{\Omega}} |\nabla v|^2, \quad v \in \mathcal{A},$$

$$\mathcal{A} := \{v \in H_\varphi^1(\Omega) \mid \nabla v \equiv 0 \text{ in } D_1 \cup D_2\}.$$

The readers can refer to the Appendix for the proofs of the above statements.

It was proved by Ammari, Kang and Lim in [7] and Ammari, Kang, H. Lee, J. Lee and Lim in [6] that, when D_1 and D_2 are balls of comparable radii embedded in $\Omega = \mathbb{R}^2$, the

blow-up rate of the gradient of the solution to the perfect conductivity problem is $\varepsilon^{-1/2}$ as ε goes to zero; with the lower bound given in [7] and the upper bound given in [6]. Yun in [31] generalized the above mentioned result in [7] by establishing the same lower bound, $\varepsilon^{-1/2}$, for two strictly convex subdomains in \mathbb{R}^2 . In this part, we give both lower and upper bounds to blow-up rates of the gradient for the solution to the perfect conductivity problem in a bounded matrix, where two strictly convex subdomains are embedded. Our methods apply to dimension $n \geq 3$ as well. One might reasonably suspect that the blow-up rate in dimension $n \geq 3$ should be smaller than that in dimension $n = 2$. However we prove the opposite: As ε goes to zero, the blow-up rate is $\varepsilon^{-1/2}$, $(\varepsilon |\ln \varepsilon|)^{-1}$ and ε^{-1} for $n = 2, 3$ and $n \geq 4$, respectively. We also give a criteria, in terms of a linear functional of the boundary data φ , for the situation where the rate of blow-up is realized. Note that [6] and [7] contain also results for $k < \infty$.

We now state more precisely what it means by saying that the boundary of a domain, say Ω , is $C^{2,\alpha}$ for $0 < \alpha < 1$: In a neighborhood of every point of $\partial\Omega$, $\partial\Omega$ is the graph of some $C^{2,\alpha}$ functions of $n - 1$ variables. We define the $C^{2,\alpha}$ norm of $\partial\Omega$, denoted as $\|\partial\Omega\|_{C^{2,\alpha}}$, as the smallest positive number $\frac{1}{a}$ such that in the $2a$ -neighborhood of every point of $\partial\Omega$, identified as 0 after a possible translation and rotation of the coordinates so that $x_n = 0$ is the tangent to $\partial\Omega$ at 0, $\partial\Omega$ is given by the graph of a $C^{2,\alpha}$ function, denoted as f , which is defined as $|x'| < a$, the a -neighborhood of 0 in the tangent plane. Moreover, $\|f\|_{C^{2,\alpha}(|x'| < a)} \leq \frac{1}{a}$.

Theorem 1.1 *Let $\Omega, D_1, D_2 \subset \mathbb{R}^n$, ε be defined as in (1.1), $\varphi \in C^2(\partial\Omega)$. Let $u \in H^1(\Omega) \cap C^1(\bar{\Omega})$ be the solution to equation (1.4). For ε sufficiently small, there is a positive constant C which depends only on $n, \kappa_0, r_0, \|\partial\Omega\|_{C^{2,\alpha}}, \|\partial D_1\|_{C^{2,\alpha}}$ and $\|\partial D_2\|_{C^{2,\alpha}}$, but independent of ε such that*

$$\begin{aligned} \|\nabla u\|_{L^\infty(\bar{\Omega})} &\leq \frac{C}{\sqrt{\varepsilon}} \|\varphi\|_{C^2(\partial\Omega)} & \text{for } n = 2, \\ \|\nabla u\|_{L^\infty(\bar{\Omega})} &\leq \frac{C}{\varepsilon |\ln \varepsilon|} \|\varphi\|_{C^2(\partial\Omega)} & \text{for } n = 3, \\ \|\nabla u\|_{L^\infty(\bar{\Omega})} &\leq \frac{C}{\varepsilon} \|\varphi\|_{C^2(\partial\Omega)} & \text{for } n \geq 4. \end{aligned} \tag{1.5}$$

Remark 1.1 *We draw attention of readers to the independent work of Yun [32] where he has also established the upper bound, $\varepsilon^{-1/2}$, in \mathbb{R}^2 . The methods are very different. Results*

in this thesis and those in [31] and [32] do not really need D_1 and D_2 to be strictly convex, the strict convexity is only needed for the portions in a fixed neighborhood (the size of the neighborhood is independent of ε) of a pair of points on ∂D_1 and ∂D_2 which realize minimal distance ε . In fact, our proofs of Theorem 1.1–1.2 also apply, with minor modification, to more general situations where two inclusions, D_1 and D_2 , are not necessarily convex near points on the boundaries where minimal distance ε is realized; see discussions after the proofs of Theorem 1.1–1.2 in Section 2.1.3.

To prove Theorem 1.1, we first decompose the solution u of equation (1.4) as follows:

$$u = C_1 v_1 + C_2 v_2 + v_3 \quad (1.6)$$

where $C_i := C_i(\varepsilon)$ ($i = 1, 2$) be the boundary value of u on ∂D_i ($i = 1, 2$) respectively, and $v_i \in C^2(\widetilde{\Omega})$ ($i = 1, 2, 3$) satisfies

$$\begin{cases} \Delta v_1 = 0 & \text{in } \widetilde{\Omega}, \\ v_1 = 1 \text{ on } \partial D_1, \quad v_1 = 0 \text{ on } \partial D_2 \cup \partial \Omega, \end{cases} \quad (1.7)$$

$$\begin{cases} \Delta v_2 = 0 & \text{in } \widetilde{\Omega}, \\ v_2 = 1 \text{ on } \partial D_2, \quad v_2 = 0 \text{ on } \partial D_1 \cup \partial \Omega, \end{cases} \quad (1.8)$$

$$\begin{cases} \Delta v_3 = 0 & \text{in } \widetilde{\Omega}, \\ v_3 = 0 \text{ on } \partial D_1 \cup \partial D_2, \quad v_3 = \varphi \text{ on } \partial \Omega. \end{cases} \quad (1.9)$$

Define

$$Q_\varepsilon[\varphi] := \int_{\partial D_1} \frac{\partial v_3}{\partial \nu} \int_{\partial \Omega} \frac{\partial v_2}{\partial \nu} - \int_{\partial D_2} \frac{\partial v_3}{\partial \nu} \int_{\partial \Omega} \frac{\partial v_1}{\partial \nu}, \quad (1.10)$$

then $Q_\varepsilon : C^2(\partial \Omega) \rightarrow \mathbb{R}$ is a linear functional.

Theorem 1.2 *With the same conditions in Theorem 1.1, let $u \in H^1(\Omega) \cap C^1(\widetilde{\Omega})$ be the solution to equation (1.4). For ε sufficiently small, there exists a positive constant C which depends on n , κ_0 , r_0 , $\|\partial \Omega\|_{C^{2,\alpha}}$, $\|\partial D_1\|_{C^{2,\alpha}}$, $\|\partial D_2\|_{C^{2,\alpha}}$ and $\|\varphi\|_{C^2(\partial \Omega)}$, but is independent of ε such that*

$$\begin{aligned} \|\nabla u\|_{L^\infty(\widetilde{\Omega})} &\geq \frac{|Q_\varepsilon[\varphi]|}{C} \cdot \frac{1}{\sqrt{\varepsilon}} & \text{for } n = 2, \\ \|\nabla u\|_{L^\infty(\widetilde{\Omega})} &\geq \frac{|Q_\varepsilon[\varphi]|}{C} \cdot \frac{1}{\varepsilon |\ln \varepsilon|} & \text{for } n = 3, \\ \|\nabla u\|_{L^\infty(\widetilde{\Omega})} &\geq \frac{|Q_\varepsilon[\varphi]|}{C} \cdot \frac{1}{\varepsilon} & \text{for } n \geq 4. \end{aligned} \quad (1.11)$$

Remark 1.2 If $\varphi \equiv 0$, then the solution to equation (1.4) is $u \equiv 0$. Theorem 1.1 and Theorem 1.2 are obvious in this case. So we only need to prove them for $\|\varphi\|_{C^2(\partial\Omega)} = 1$, by considering $u/\|\varphi\|_{C^2(\partial\Omega)}$.

Remark 1.3 It is interesting to know when $|Q_\varepsilon[\varphi]| \geq \frac{1}{C}$ for some positive constant C independent of ε . Roughly speaking $Q_\varepsilon[\varphi] \rightarrow Q^*[\varphi]$ as $\varepsilon \rightarrow 0$, and this amounts to $Q^*[\varphi] \neq 0$. For details, see Section 2.2.

Theorem 1.1–1.2 can be extended to equations with more general coefficients as follows: Let $n, \Omega, D_1, D_2, \varepsilon$ and φ be same as in Theorem 1.1, and let

$$A_2(x) := (a_2^{ij}(x)) \in C^2(\overline{\Omega})$$

be $n \times n$ symmetric matrix functions in $\tilde{\Omega}$ satisfying for some constants $0 < \lambda \leq \Lambda < \infty$,

$$\lambda|\xi|^2 \leq a_2^{ij}(x)\xi_i\xi_j \leq \Lambda|\xi|^2, \quad \forall x \in \tilde{\Omega}, \quad \forall \xi \in \mathbb{R}^n, \quad (1.12)$$

and $a_2^{ij}(x) \in C^2(\overline{\Omega \setminus D_1 \cup D_2})$.

We consider

$$\left\{ \begin{array}{ll} \partial_{x_j} (a_2^{ij}(x) \partial_{x_i} u) = 0 & \text{in } \tilde{\Omega}, \\ u|_+ = u|_- & \text{on } \partial D_1 \cup \partial D_2, \\ \nabla u = 0 & \text{in } D_1 \cup D_2, \\ \int_{\partial D_i} a_2^{ij}(x) \partial_{x_i} u \nu_j|_+ = 0 & (i = 1, 2), \\ u = \varphi & \text{on } \partial\Omega. \end{array} \right. \quad (1.13)$$

where repeated indices denote as usual summations.

Here is an extension of Theorem 1.1:

Theorem 1.3 With the above assumptions, let $u \in H^1(\Omega) \cap C^1(\overline{\tilde{\Omega}})$ be the solution to equation (1.13). For ε sufficient small, there is a positive constant C which depends only on $n, \kappa_0, r_0, \|\partial\Omega\|_{C^{2,\alpha}}, \|\partial D_1\|_{C^{2,\alpha}}, \|\partial D_2\|_{C^{2,\alpha}}, \lambda, \Lambda$ and $\|A_2\|_{C^2(\overline{\tilde{\Omega}})}$, but independent of ε such that estimate (1.5) holds.

Similar to the decomposition formula (1.6), we decompose the solution u of equation (1.13) as follows:

$$u = C_1 V_1 + C_2 V_2 + V_3 \quad (1.14)$$

where $C_i := C_i(\varepsilon)$ ($i = 1, 2$) be the boundary value of u on ∂D_i ($i = 1, 2$) respectively, and $V_i \in C^2(\overline{\Omega})$ ($i = 1, 2, 3$) satisfies

$$\begin{cases} \partial_{x_j} \left(a_2^{ij}(x) \partial_{x_i} V_1 \right) = 0 & \text{in } \tilde{\Omega}, \\ V_1 = 1 \text{ on } \partial D_1, \quad V_1 = 0 \text{ on } \partial D_2 \cup \partial \Omega, \end{cases} \quad (1.15)$$

$$\begin{cases} \partial_{x_j} \left(a_2^{ij}(x) \partial_{x_i} V_2 \right) = 0 & \text{in } \tilde{\Omega}, \\ V_2 = 1 \text{ on } \partial D_2, \quad V_2 = 0 \text{ on } \partial D_1 \cup \partial \Omega, \end{cases} \quad (1.16)$$

$$\begin{cases} \partial_{x_j} \left(a_2^{ij}(x) \partial_{x_i} V_3 \right) = 0 & \text{in } \tilde{\Omega}, \\ V_3 = 0 \text{ on } \partial D_1 \cup \partial D_2, \quad V_3 = \varphi \text{ on } \partial \Omega. \end{cases} \quad (1.17)$$

Define

$$\begin{aligned} Q_\varepsilon[\varphi] := & \int_{\partial D_1} a_2^{ij}(x) \partial_{x_i} V_3 \nu_j \int_{\partial \Omega} a_2^{ij}(x) \partial_{x_i} V_2 \nu_j \\ & - \int_{\partial D_2} a_2^{ij}(x) \partial_{x_i} V_3 \nu_j \int_{\partial \Omega} a_2^{ij}(x) \partial_{x_i} V_1 \nu_j, \end{aligned} \quad (1.18)$$

then $Q_\varepsilon : C^2(\partial \Omega) \rightarrow \mathbb{R}$ is a linear functional.

Theorem 1.4 *With the same conditions in Theorem 1.3, let $u \in H^1(\Omega) \cap C^1(\overline{\Omega})$ be the solution to equation (1.13). For ε sufficiently small and $Q_\varepsilon[\varphi]$ defined by (1.18), there is a positive constant C which depends only on $n, \kappa_0, r_0, \|\partial D_1\|_{C^{2,\alpha}}, \|\partial D_2\|_{C^{2,\alpha}}, \lambda, \Lambda$ and $\|A_2\|_{C^2(\overline{\Omega})}$, but independent of ε such that estimate (1.11) holds.*

Part II. Along the approach in the first part, we have extended Theorem 1.1 and 1.3 from two inclusions to multiple inclusions, see Theorem 1.5 below.

The complementary problem to the perfect case is the insulated case when $k = 0$ in (1.2). For $n = 2$, Ammari, Kang, H. Lee, J. Lee and Lim have given in [7] and [6] the optimal bound when D_1 and D_2 are balls of comparable radii embedded in $\Omega = \mathbb{R}^2$. The blow-up rate of the gradient of solutions is $\varepsilon^{-1/2}$ as the distance between D_1 and D_2 , ε ,

goes to zero. They obtained this by converting the insulated case to the perfect case using harmonic conjugators. The situation for $n \geq 3$ is different since the $k = 0$ case can not be converted to the $k = \infty$ case. In this part, we proved that the above mentioned optimal upper bound $\epsilon^{-1/2}$ is also an upper bound of $\|\nabla u\|_{L^\infty}$ for $n \geq 3$. In fact, what we have obtained is a local version of the estimates, see Theorem 1.6 below. On the other hand, we do not know yet whether the estimates are sharp for $n \geq 3$.

Now let, as before, Ω be a bounded open set in \mathbb{R}^n with $C^{2,\alpha}$ boundary, $n \geq 2$, $0 < \alpha < 1$. Instead of two inclusions, let $\{D_i\}, i = 1, 2, \dots, m$, be m strictly convex open subsets in Ω with $C^{2,\alpha}$ boundaries, $m \geq 2$, satisfying

$$\begin{aligned} \overline{D_i} &\subset \Omega, \quad \text{the principal curvature of } \partial D_i \geq \kappa_0, \\ \varepsilon_{ij} &:= \text{dist}(D_i, D_j) > 0, \quad (i \neq j) \\ \text{dist}(D_i, \partial\Omega) &> r_0, \quad \text{diam}(\Omega) < \frac{1}{r_0}, \end{aligned} \tag{1.19}$$

where $\kappa_0, r_0 > 0$ are universal constants independent of ε_{ij} . As before we will assume that the $C^{2,\alpha}$ norms of ∂D_i are under control, while ε_{ij} can become very small.

For $i \neq j$, let

$$\text{dist}(x_{ij}^i, x_{ij}^j) = \text{dist}(D_i, D_j) = \varepsilon_{ij} > 0, \quad x_{ij}^i \in \partial D_i, \quad x_{ij}^j \in \partial D_j,$$

and

$$x_{ij}^0 := \frac{1}{2}(x_{ij}^i + x_{ij}^j).$$

It is easy to see that there exists some positive constant δ which depends only on κ_0, r_0 and $\{\|\partial D_i\|_{C^{2,\alpha}}\}$, but is independent of $\{\varepsilon_{ij}\}$ such that any ball of radius 2δ can intersect at most two elements in $\{D_i\}$. We will only be interested in those pair i, j satisfying

$$B_{ij} := B(x_{ij}^0, \delta) \text{ intersects both } D_i \text{ and } D_j. \tag{1.20}$$

Given $\varphi \in C^2(\partial\Omega)$, consider, for $m \geq 2$,

$$\begin{cases} \partial_{x_j} \left(a_2^{ij}(x) \partial_{x_i} u \right) = 0 & \text{in } \tilde{\Omega} := \Omega \setminus \overline{\bigcup_{i=1}^m D_i}, \\ u|_+ = u|_- & \text{on } \partial D_i \ (i = 1, 2, \dots, m), \\ \nabla u = 0 & \text{on } D_i \ (i = 1, 2, \dots, m), \\ \int_{\partial D_i} a_2^{ij}(x) \partial_{x_i} u \nu_j|_+ = 0 & (i = 1, 2, \dots, m), \\ u = \varphi & \text{on } \partial\Omega, \end{cases} \quad (1.21)$$

where $a_2^{ij}(x)$ satisfies condition (1.12). Then we have

Theorem 1.5 *Let $\Omega, \{D_l\} \subset \mathbb{R}^n$, $\{\varepsilon_{kl}\}$ be defined as in (1.19), $n \geq 2$, $\varphi \in C^2(\partial\Omega)$, and let $u \in H^1(\Omega) \cap C^1(\tilde{\Omega})$ be the solution to equation (1.21), with $u \equiv C_l$ on D_l . Then for any pair i, j satisfying (1.20),*

$$\begin{aligned} \|\nabla u\|_{L^\infty(\tilde{\Omega} \cap B_{ij})} &\leq C \frac{|C_i - C_j|}{\varepsilon_{ij}} + C \|\varphi\|_{C^2(\partial\Omega)} \\ &\leq \begin{cases} \frac{C}{\sqrt{\varepsilon_{ij}}} \|\varphi\|_{C^2(\partial\Omega)} & \text{for } n = 2, \\ \frac{C}{\varepsilon_{ij} |\ln \varepsilon_{ij}|} \|\varphi\|_{C^2(\partial\Omega)} & \text{for } n = 3, \\ \frac{C}{\varepsilon_{ij}} \|\varphi\|_{C^2(\partial\Omega)} & \text{for } n \geq 4, \end{cases} \end{aligned} \quad (1.22)$$

where C depends only on $n, m, \kappa_0, r_0, \|\partial\Omega\|_{C^{2,\alpha}}, \{\|\partial D_l\|_{C^{2,\alpha}}\}, \lambda, \Lambda$ and $\{\|a_2^{kl}\|_{C^2(\bar{\Omega})}\}$.

All our previous theorems concern perfect inclusions ($k = \infty$). Now we consider the complementary problem when $k = 0$ (insulated inclusions) in (1.2).

Let $\Omega, D_i \subset \mathbb{R}^n$, ε_{ij} be defined as in (1.19), $\varphi \in C^2(\partial\Omega)$. The insulated conductivity problem can be described as follows, for $i = 1, 2, \dots, m$,

$$\begin{cases} \partial_{x_j} \left(a_2^{ij}(x) \partial_{x_i} u \right) = 0 & \text{in } \tilde{\Omega}, \\ a_2^{ij}(x) \partial_{x_i} u \nu_j|_+ = 0 & \text{on } \partial D_i, \\ u = \varphi & \text{on } \partial\Omega. \end{cases} \quad (1.23)$$

The existence and uniqueness of solutions to equation (1.23) are elementary. By the maximum principle, $\|u\|_{L^\infty(\tilde{\Omega})} \leq \|\varphi\|_{L^\infty(\partial\Omega)}$.

To obtain an upper bound of $\|\nabla u\|_{L^\infty(\tilde{\Omega})}$ for solutions, we only need to consider the following local situation: For any pair i, j satisfying (1.20), consider

$$\begin{cases} \partial_{x_j} \left(a_2^{ij}(x) \partial_{x_i} u \right) = 0 & \text{in } \tilde{\Omega} \cap B(x_{ij}^0, 2\delta), \\ a_2^{ij}(x) \partial_{x_i} u \nu_j|_+ = 0 & \text{on } (\partial D_i \cup \partial D_j) \cap B(x_{ij}^0, 2\delta), \\ |u| \leq 1 & \text{in } \tilde{\Omega} \cap B(x_{ij}^0, 2\delta). \end{cases} \quad (1.24)$$

Then we have

Theorem 1.6 *Let $\Omega, \{D_l\} \subset \mathbb{R}^n$, $\{\varepsilon_{kl}\}$ be defined as in (1.19), $n \geq 2$. For any pair i, j satisfying (1.20), let $u \in C^1(\overline{\tilde{\Omega} \cap B(x_{ij}^0, 2\delta)})$ be a solution to equation (1.24). Then*

$$\|\nabla u\|_{L^\infty(\tilde{\Omega} \cap B_{ij})} \leq \frac{C}{\sqrt{\varepsilon_{ij}}}, \quad (1.25)$$

where the constant C depends only on $n, m, \kappa_0, r_0, \|\partial D_i\|_{C^{2,\alpha}}, \|\partial D_j\|_{C^{2,\alpha}}, \lambda, \Lambda$ and $\{\|a_2^{kl}\|_{C^2(\bar{\Omega})}\}$.

Chapter 2

The perfect conductivity problem with two inclusions

This chapter concerns the gradient estimates of the perfect conductivity problem with two inclusions. We will prove Theorem 1.1–1.4. In Section 2.1 we first prove Theorem 1.1–1.2. In Section 2.2 we give a criteria for $|Q_\varepsilon[\varphi]|$ to be bounded below by a positive constant independent of ε . Theorem 1.3–1.4 are proved in Section 2.3. In the Appendix A we present some elementary results for the perfect conductivity problem.

2.1 Proof of Theorem 1.1 and 1.2

In the introduction, we write $u = C_1 v_1 + C_2 v_2 + v_3$ as in (1.6). To prove our main theorems, we first estimate $\|\nabla u\|_{L^\infty(\tilde{\Omega})}$ in terms of $|C_1 - C_2|$, and then estimate $|C_1 - C_2|$.

In this section we use, unless otherwise stated, C to denote various positive constants whose values may change from line to line and which depend only on $n, \kappa_0, r_0, \|\partial\Omega\|_{C^{2,\alpha}}, \|\partial D_1\|_{C^{2,\alpha}}$ and $\|\partial D_2\|_{C^{2,\alpha}}$.

Proposition 2.1 *Under the hypotheses of Theorem 1.1, let u be the solution of equation (1.4). There exists a positive constants C , such that, for sufficiently small $\varepsilon > 0$,*

$$\frac{1}{\varepsilon} |C_1 - C_2| \leq \|\nabla u\|_{L^\infty(\tilde{\Omega})} \leq \frac{C}{\varepsilon} |C_1 - C_2| + C\|\varphi\|_{C^2(\partial\Omega)}. \quad (2.1)$$

To prove this proposition, we first estimate the gradients of v_1, v_2 and v_3 . Without loss of generality, we may assume throughout the proof of the proposition that $\|\varphi\|_{C^2(\partial\Omega)} = 1$; see Remark 1.2.

Lemma 2.1 *Let v_1, v_2 be defined by equations (1.7) and (1.8), then for $n \geq 2$, we have*

$$\|\nabla v_1\|_{L^\infty(\tilde{\Omega})} + \|\nabla v_2\|_{L^\infty(\tilde{\Omega})} \leq \frac{C}{\varepsilon}, \quad \left\| \frac{\partial v_1}{\partial \nu} \right\|_{L^\infty(\partial\Omega)} + \left\| \frac{\partial v_2}{\partial \nu} \right\|_{L^\infty(\partial\Omega)} \leq C.$$

Proof: By the maximum principle, $\|v_1\|_{L^\infty(\tilde{\Omega})} \leq 1$, and since v_1 achieves constants on each connected component of $\partial\tilde{\Omega}$, and each connected component of $\partial\tilde{\Omega}$ is $C^{2,\alpha}$ then the gradient estimates for harmonic functions implies that

$$\|\nabla v_1\|_{L^\infty(\tilde{\Omega})} \leq \frac{C\|v_1\|_{L^\infty}}{\text{dist}(\partial D_1, \partial D_2)} = \frac{C}{\varepsilon}.$$

Similarly, we can prove $\|\nabla v_2\|_{L^\infty(\tilde{\Omega})} \leq C/\varepsilon$. The second inequality follows from the boundary estimates for harmonic functions. \square

Before estimating $|\nabla v_3|$, we first prove:

Lemma 2.2 *Let $\rho \in C^2(\tilde{\Omega})$ be the solution to:*

$$\begin{cases} \Delta \rho = 0 & \text{in } \tilde{\Omega}, \\ \rho = 0 \text{ on } \partial D_1 \cup \partial D_2, & \rho = 1 \text{ on } \partial \Omega. \end{cases} \quad (2.2)$$

Then $\|\nabla \rho\|_{L^\infty(\tilde{\Omega})} \leq C$.

Proof: Let $\rho_i (i = 1, 2) \in C^2(\Omega \setminus \overline{D_i}) \cap C^1(\overline{\Omega \setminus D_i})$ be the solution to:

$$\begin{cases} \Delta \rho_i = 0 & \text{in } \Omega \setminus \overline{D_i}, \\ \rho_i = 0 \text{ on } \partial D_i, & \rho_i = 1 \text{ on } \partial \Omega. \end{cases}$$

Again by the maximum principle and the strong maximum principle, we obtain $0 < \rho_1 < 1$ in $\Omega \setminus \overline{D_1}$. Since $\overline{D_2} \subset \Omega \setminus \overline{D_1}$, we have $\rho_1 > 0 = \rho$ on ∂D_2 . And since $\rho_1 = \rho$ on ∂D_1 and $\partial \Omega$, therefore $\rho_1 > \rho$ on $\tilde{\Omega}$. Now because $\rho_1 = \rho = 0$ on ∂D_1 and $\rho_1 > \rho > 0$ on $\tilde{\Omega}$, so

$$\|\nabla \rho\|_{L^\infty(\partial D_1)} \leq \|\nabla \rho_1\|_{L^\infty(\partial D_1)} \leq C.$$

Similarly,

$$\|\nabla \rho\|_{L^\infty(\partial D_2)} \leq \|\nabla \rho_2\|_{L^\infty(\partial D_2)} \leq C.$$

By the boundary estimate of harmonic functions, we know that $\|\nabla \rho\|_{L^\infty(\partial \Omega)} \leq C$.

Since $\Delta \rho = 0$ in $\tilde{\Omega}$, $\partial_{x_i} \rho$ is also harmonic, by the maximum principle,

$$\|\nabla \rho\|_{L^\infty(\tilde{\Omega})} \leq \max \left(\|\nabla \rho\|_{L^\infty(\partial D_1)}, \|\nabla \rho\|_{L^\infty(\partial D_2)}, \|\nabla \rho\|_{L^\infty(\partial \Omega)} \right) \leq C.$$

\square

Now, we estimate $|\nabla v_3|$:

Lemma 2.3 *Let v_3 be defined by equation (1.9), for $n \geq 2$, we have*

$$\|\nabla v_3\|_{L^\infty(\tilde{\Omega})} \leq C.$$

Proof: Since $v_3 = -\rho = \rho = 0$ on $\partial D_i (i = 1, 2)$, and $-\rho \leq v_3 = \varphi \leq \rho$ on $\partial\Omega$, we have, by the maximum principle,

$$-\rho \leq v_3 \leq \rho \quad \text{in } \tilde{\Omega}.$$

It follows, for $i = 1, 2$, that

$$\|\nabla v_3\|_{L^\infty(\partial D_i)} \leq \|\nabla \rho\|_{L^\infty(\partial D_i)} \leq C.$$

By the boundary estimate,

$$\|\nabla v_3\|_{L^\infty(\partial\Omega)} \leq C.$$

By the harmonicity of $\partial_{x_i} v_3$ and the maximum principle,

$$\|\nabla v_3\|_{L^\infty(\tilde{\Omega})} \leq C.$$

□

Remark 2.1 *Without assuming $\|\varphi\|_{C^2(\partial\Omega)} = 1$, we have*

$$\|\nabla v_3\|_{L^\infty(\partial D_1 \cup \partial D_2)} \leq C \|\varphi\|_{L^\infty(\partial\Omega)},$$

where C has the dependence specified at the beginning of this section, except that it does not depend on $\|\partial\Omega\|_{C^{2,\alpha}}$. This is easy to see from the proof of Lemma 2.3.

The above lemma yields the main result of [2].

Corollary 2.1 ([2]) *Let B_1 and B_2 be two spheres with radius R and centered at $(\pm R \pm \frac{\varepsilon}{2}, 0, \dots, 0)$, respectively. Let H be a harmonic function in \mathbb{R}^3 . Define u to be the solution to*

$$\begin{cases} \Delta u = 0 & \text{in } \mathbb{R}^3 \setminus \overline{B_1 \cup B_2}, \\ u = 0 & \text{on } \partial B_1 \cup \partial B_2, \\ u(x) - H(x) = O(|x|^{-1}) & \text{as } |x| \rightarrow +\infty. \end{cases}$$

Then there is a constant C independent of ε such that

$$\|\nabla(u - H)\|_{L^\infty(\mathbb{R}^3 \setminus \overline{B_1 \cup B_2})} \leq C.$$

Proof: By the maximum principle and interior estimates of harmonic functions, the C^3 norm of $u|_{B_{2R}(0)}$ is bounded by a constant independent of ε . Apply Lemma 2.3 with $\Omega = B_{2R}(0)$ and $\varphi = u|_{B_{2R}(0)}$, we immediately obtain the above corollary. \square

With the above lemmas, we give the

Proof of Proposition 2.1: Since $u = C_1$ on ∂D_1 , $u = C_2$ on ∂D_2 , and $\text{dist}(\partial D_1, \partial D_2) = \varepsilon$, by the mean value theorem, $\exists \xi \in \tilde{\Omega}$ such that

$$\|\nabla u\|_{L^\infty(\tilde{\Omega})} \geq |\nabla u(\xi)| \geq \frac{|C_1 - C_2|}{\varepsilon}.$$

By the decomposition formula (1.6),

$$\nabla u = C_1 \nabla v_1 + C_2 \nabla v_2 + \nabla v_3 = (C_1 - C_2) \nabla v_1 + C_2 \nabla(v_1 + v_2) + \nabla v_3.$$

Hence,

$$\|\nabla u\|_{L^\infty(\tilde{\Omega})} \leq |C_1 - C_2| \|\nabla v_1\|_{L^\infty(\tilde{\Omega})} + |C_2| \|\nabla(v_1 + v_2)\|_{L^\infty(\tilde{\Omega})} + \|\nabla v_3\|_{L^\infty(\tilde{\Omega})}.$$

By Lemma 2.2, since $v_1 + v_2 = 1 - \rho$ in $\tilde{\Omega}$, we have

$$\|\nabla(v_1 + v_2)\|_{L^\infty(\tilde{\Omega})} = \|\nabla(1 - \rho)\|_{L^\infty(\tilde{\Omega})} = \|\nabla \rho\|_{L^\infty(\tilde{\Omega})} \leq C.$$

Using the fact we showed in the Appendix, $\|u\|_{H^1(\Omega)} \leq C$, so $|C_1| + |C_2| \leq C$.

Therefore using also Lemma 2.1 we obtain,

$$\|\nabla u\|_{L^\infty(\tilde{\Omega})} \leq \frac{C}{\varepsilon} |C_1 - C_2| + C.$$

This proof is now completed. \square

Later we will give an estimate of $|C_1 - C_2|$, which, together with Proposition 2.1, yields the lower and upper bounds of $\|\nabla u\|_{L^\infty(\tilde{\Omega})}$ for strictly convex subdomains D_1 and D_2 .

2.1.1 Estimate of $|C_1 - C_2|$

Back to the decomposition formula (1.6), denote

$$a_{ij} = \int_{\partial D_i} \frac{\partial v_j}{\partial \nu} \quad (i, j = 1, 2), \quad b_i = \int_{\partial D_i} \frac{\partial v_3}{\partial \nu} \quad (i = 1, 2). \quad (2.3)$$

We first give some basic lemmas:

Lemma 2.4 *Let a_{ij} and b_i be defined as in (2.3), then they satisfy the following:*

1. $a_{12} = a_{21} > 0$, $a_{11} < 0$, $a_{22} < 0$,
2. $-C \leq a_{11} + a_{21} \leq -\frac{1}{C}$, $-C \leq a_{22} + a_{12} \leq -\frac{1}{C}$,
3. $|b_1| \leq C$, $|b_2| \leq C$.

By the fourth line of equation (1.4), C_1 and C_2 satisfy

$$\begin{cases} a_{11}C_1 + a_{12}C_2 + b_1 = 0, \\ a_{21}C_1 + a_{22}C_2 + b_2 = 0. \end{cases} \quad (2.4)$$

By solving the above linear system, using $a_{12} = a_{21}$ and $a_{11}a_{22} - a_{12}a_{21} > 0$ which follows from Lemma 2.4, we obtain

$$C_1 = \frac{-b_1a_{22} + b_2a_{12}}{a_{11}a_{22} - a_{12}^2}, \quad C_2 = \frac{-b_2a_{11} + b_1a_{12}}{a_{11}a_{22} - a_{12}^2}, \quad (2.5)$$

and therefore,

$$|C_1 - C_2| = \frac{|b_1 - \alpha b_2|}{|a_{11} - \alpha a_{12}|}, \quad \text{where } \alpha = \frac{a_{11} + a_{12}}{a_{22} + a_{12}} > 0. \quad (2.6)$$

Based on this formula, we will give the estimates for $|a_{11} - \alpha a_{12}|$ and $|b_1 - \alpha b_2|$, then the estimate for $|C_1 - C_2|$ follows immediately.

Proof of Lemma 2.4: (1) By the maximum principle and the strong maximum principle,

$$0 < v_1 < 1 \quad \text{in } \tilde{\Omega}.$$

By the Hopf Lemma, we know that

$$\frac{\partial v_1}{\partial \nu} \Big|_{\partial D_1} < 0, \quad \frac{\partial v_1}{\partial \nu} \Big|_{\partial D_2} > 0, \quad \frac{\partial v_1}{\partial \nu} \Big|_{\partial \Omega} < 0.$$

Similarly,

$$\frac{\partial v_2}{\partial \nu}|_{\partial D_1} > 0, \quad \frac{\partial v_2}{\partial \nu}|_{\partial D_2} < 0, \quad \frac{\partial v_2}{\partial \nu}|_{\partial \Omega} < 0.$$

Thus $a_{11} < 0$, $a_{12} > 0$, $a_{21} > 0$ and $a_{22} < 0$.

Also, since v_1 and v_2 are the solutions of equations (1.7) and equations (1.8), respectively, we have

$$\begin{aligned} 0 &= \int_{\tilde{\Omega}} \Delta v_1 \cdot v_2 - \int_{\tilde{\Omega}} \Delta v_2 \cdot v_1 = - \int_{\partial D_2} \frac{\partial v_1}{\partial \nu} \cdot 1 + \int_{\partial D_1} \frac{\partial v_2}{\partial \nu} \cdot 1 \\ &= -a_{21} + a_{12}, \end{aligned} \tag{2.7}$$

i.e. $a_{21} = a_{12}$.

(2) We will prove the first inequality, the second one stands with the same reason. By the harmonicity of v_1 in $\tilde{\Omega}$,

$$a_{11} + a_{21} = - \int_{\tilde{\Omega}} \Delta v_1 + \int_{\partial \Omega} \frac{\partial v_1}{\partial \nu} = \int_{\partial \Omega} \frac{\partial v_1}{\partial \nu} < 0.$$

By Lemma 2.1,

$$a_{11} + a_{21} = \int_{\partial \Omega} \frac{\partial v_1}{\partial \nu} \geq -C.$$

On the other hand, since $0 < v_1 < 1$ in $\tilde{\Omega}$ and $v_1 = 1$ on ∂D_1 , by the boundary gradient estimates of a harmonic function, $\exists B(\bar{x}, 2\bar{r}) \subset \tilde{\Omega}$, such that $v_1 > 1/2$ in $B(\bar{x}, \bar{r})$, where \bar{r} is independent of ε . Let $\rho \in C^2(\Omega \setminus \overline{D_2 \cup B(\bar{x}, \bar{r})}) \cup C^1(\partial \Omega \cup \partial D_2 \cup \partial B(\bar{x}, \bar{r}))$ be the solution of the following equation:

$$\begin{cases} \Delta \rho = 0 & \text{in } \Omega \setminus \overline{D_2 \cup B(\bar{x}, \bar{r})}, \\ \rho = 1/2 \text{ on } \partial B(\bar{x}, \bar{r}) & \rho = 0 \text{ on } \partial D_2 \cup \partial \Omega. \end{cases}$$

By the maximum principle and the strong maximum principle, $0 < \rho < 1/2$ in $\Omega \setminus \overline{D_2 \cup B(\bar{x}, \bar{r})}$.

A contradiction argument based on the Hopf Lemma yields,

$$-\frac{\partial \rho}{\partial \nu} \geq \frac{1}{C} \quad \text{on } \partial \Omega.$$

On the other hand, since $\rho \leq v_1$ on the boundary of $\Omega \setminus \overline{D_1 \cup D_2 \cup B(\bar{x}, \bar{r})}$, we obtain, via the maximum principle, $0 < \rho \leq v_1$ in $\Omega \setminus \overline{D_1 \cup D_2 \cup B(\bar{x}, \bar{r})}$. It follows, using $\rho = v_1 = 0$ on $\partial \Omega$, that

$$\frac{\partial v_1}{\partial \nu} \leq \frac{\partial \rho}{\partial \nu} \quad \text{on } \partial \Omega.$$

Thus,

$$a_{11} + a_{21} = \int_{\partial\Omega} \frac{\partial v_1}{\partial \nu} \leq \int_{\partial\Omega} \frac{\partial \rho}{\partial \nu} \leq -\frac{1}{C}.$$

(3) Clearly,

$$0 = \int_{\tilde{\Omega}} \Delta v_1 \cdot v_3 - \int_{\tilde{\Omega}} \Delta v_3 \cdot v_1 = \int_{\partial\Omega} \frac{\partial v_1}{\partial \nu} \cdot \varphi + \int_{\partial D_1} \frac{\partial v_3}{\partial \nu} \cdot 1 = \int_{\partial\Omega} \frac{\partial v_1}{\partial \nu} \cdot \varphi + b_1.$$

Thus,

$$|b_1| = \left| \int_{\partial\Omega} \frac{\partial v_1}{\partial \nu} \cdot \varphi \right| \leq \int_{\partial\Omega} \left| \frac{\partial v_1}{\partial \nu} \right| \leq C.$$

Therefore, we finished the proof. \square

2.1.2 Estimate of $|a_{11} - \alpha a_{12}|$

By a translation and rotation of the axis, we may assume without loss of generality that D_1, D_2 are two strictly convex subdomains in $\Omega \subset \mathbb{R}^n$ which satisfy the following:

$$(-\varepsilon/2, 0') \in \partial D_1, \quad (\varepsilon/2, 0') \in \partial D_2, \quad \varepsilon = \text{dist}(D_1, D_2). \quad (2.8)$$

Near the origin, we can find a ball $B(0, r)$ such that the portion of ∂D_i ($i = 1, 2$) in $B(0, r)$ is strictly convex, where $r > 0$ is independent of ε . Then $\partial D_1 \cap B(0, r)$ and $\partial D_2 \cap B(0, r)$ can be represented by the graph of $x_1 = f(x') - \varepsilon/2$ and $x_1 = g(x') + \varepsilon/2$ respectively, where $x' = (x_2, \dots, x_n)$. Thus $f(0') = g(0') = 0$, $\nabla f(0') = \nabla g(0') = 0$, and $-CI \leq (D^2 f(0')) \leq -\frac{1}{C}I$, $\frac{1}{C}I \leq (D^2 g(0')) \leq CI$.

With these notations, we first estimate a_{ii} for $i = 1, 2$.

Lemma 2.5 *Let a_{ii} be defined by (2.3), then*

$$\frac{1}{C\sqrt{\varepsilon}} \leq -a_{ii} \leq \frac{C}{\sqrt{\varepsilon}}, \quad \text{for } n = 2, \quad i = 1, 2.$$

Proof: It suffices to prove it for a_{11} . By the harmonicity of v_1 , we have

$$0 = \int_{\tilde{\Omega}} \Delta v_1 \cdot v_1 = - \int_{\tilde{\Omega}} |\nabla v_1|^2 - \int_{\partial D_1} \frac{\partial v_1}{\partial \nu} = - \int_{\tilde{\Omega}} |\nabla v_1|^2 - a_{11},$$

i.e.

$$a_{11} = - \int_{\tilde{\Omega}} |\nabla v_1|^2.$$

Now we construct a function (here in \mathbb{R}^2 , we let $x = x_1$, $y = x_2$)

$$\overline{w}(x, y) = - \frac{x - g(y) - \frac{\varepsilon}{2}}{g(y) - f(y) + \varepsilon} \quad (2.9)$$

on $O_r := \tilde{\Omega} \cap \{(x, y) \mid |y| < r\}$. It is clear that $\overline{w}(x, y)$ is linear in x for fixed y and

$$\overline{w} \mid_{B(0, r) \cap \partial D_1} = 1; \quad \overline{w} \mid_{B(0, r) \cap \partial D_2} = 0,$$

so we have

$$\int_{f(y) - \frac{\varepsilon}{2}}^{g(y) + \frac{\varepsilon}{2}} |\partial_x \overline{w}(x, y)|^2 dx \leq \int_{f(y) - \frac{\varepsilon}{2}}^{g(y) + \frac{\varepsilon}{2}} |\partial_x v_1(x, y)|^2 dx,$$

i.e.

$$\frac{1}{g(y) - f(y) + \varepsilon} \leq \int_{f(y) - \frac{\varepsilon}{2}}^{g(y) + \frac{\varepsilon}{2}} |\partial_x v_1(x, y)|^2.$$

Integrating on y we get

$$\begin{aligned} \int_0^{r/2} \int_{f(y) - \frac{\varepsilon}{2}}^{g(y) + \frac{\varepsilon}{2}} |\partial_x v_1(x, y)|^2 dx dy &\geq \int_0^{r/2} \frac{1}{g(y) - f(y) + \varepsilon} dy \\ &\geq \frac{1}{C} \int_0^{r/2} \frac{1}{y^2 + \varepsilon} dy = \frac{1}{C\sqrt{\varepsilon}}. \end{aligned} \quad (2.10)$$

Thus

$$-a_{11} \geq \int_0^{r/2} \int_{f(y) - \frac{\varepsilon}{2}}^{g(y) + \frac{\varepsilon}{2}} |\partial_x v_1(x, y)|^2 dx dy \geq \frac{1}{C\sqrt{\varepsilon}}.$$

On the other hand, we can find $\psi \in C^2(\overline{\Omega})$ such that

$$\begin{aligned} \psi &= 0 \text{ on } \overline{O}_{r/8}, \quad \psi = 1 \text{ on } \partial D_1 \setminus (\overline{O}_{r/4}), \quad \psi = 0 \text{ on } \partial D_2 \setminus (\overline{O}_{r/4}), \\ \psi &= 0 \text{ on } \partial \Omega, \quad \text{and} \quad \|\nabla \psi\|_{L^\infty(\Omega)} \leq C. \end{aligned}$$

We can also find $\rho \in C^2(\overline{\Omega})$ such that

$$0 \leq \rho \leq 1, \quad \rho = 1 \text{ on } \overline{O}_{r/2}, \quad \rho = 0 \text{ on } \overline{\Omega} \setminus O_r \text{ and } |\nabla \rho| \leq C.$$

Let $w = \rho \overline{w} + (1 - \rho) \psi$, then $w = 1 = v_1$ on ∂D_1 ; $w = 0 = v_1$ on ∂D_2 ; $w = 0 = v_1$ on $\partial \Omega$ and $w = \overline{w}$ on $\overline{O}_{r/2}$. Then by the properties of ψ , ρ and the harmonicity of v_1 , we have

$$\int_{\tilde{\Omega}} |\nabla v_1|^2 \leq \int_{\tilde{\Omega}} |\nabla w|^2 \leq \int_{\tilde{\Omega} \cap O_{r/2}} |\nabla \overline{w}|^2 + C. \quad (2.11)$$

A calculation gives

$$\partial_y \bar{w} = \frac{g'(y)(g(y) - f(y) + \varepsilon) - (g(y) - x + \frac{\varepsilon}{2})(g'(y) - f'(y))}{(g(y) - f(y) + \varepsilon)^2}.$$

We will show $\int_{\tilde{\Omega} \cap O_{r/2}} |\partial_y \bar{w}|^2 \leq C$.

Indeed,

$$\begin{aligned} & \int_0^{r/2} \int_{f(y)-\frac{\varepsilon}{2}}^{g(y)+\frac{\varepsilon}{2}} |\partial_y \bar{w}(x, y)|^2 dx dy \\ & \leq 2 \int_0^{r/2} \int_{f(y)-\frac{\varepsilon}{2}}^{g(y)+\frac{\varepsilon}{2}} \left(\frac{g'(y)^2}{(g(y) - f(y) + \varepsilon)^2} + \frac{(g(y) - x + \frac{\varepsilon}{2})^2 (g'(y) - f'(y))^2}{(g(y) - f(y) + \varepsilon)^4} \right) dx dy \\ & = 2 \int_0^{r/2} \frac{g'(y)^2}{g(y) - f(y) + \varepsilon} dy + 2 \int_0^{r/2} \frac{(g'(y) - f'(y))^2}{g(y) - f(y) + \varepsilon} dy \\ & \leq C \int_0^{r/2} \frac{y^2}{y^2 + \varepsilon} dy + C \int_0^{r/2} \frac{y^2}{y^2 + \varepsilon} dy \\ & \leq C. \end{aligned} \tag{2.12}$$

Then by (2.11) and (2.12)

$$\begin{aligned} |a_{11}| &= \int_{\tilde{\Omega}} |\nabla v_1|^2 \leq \int_{\tilde{\Omega} \cap O_{r/2}} |\nabla \bar{w}|^2 + C \\ &\leq C \int_0^{r/2} \int_{f(y)-\frac{\varepsilon}{2}}^{g(y)+\frac{\varepsilon}{2}} |D_x \bar{w}(x, y)|^2 dx dy + C \\ &= C \int_0^{r/2} \frac{1}{g(y) - f(y) + \varepsilon} dy + C \leq C \int_0^{r/2} \frac{1}{y^2 + \varepsilon} dy + C \\ &\leq \frac{C}{\sqrt{\varepsilon}}. \end{aligned} \tag{2.13}$$

The proof is completed. \square

Similarly, we have

Lemma 2.6 *Let a_{ii} be defined by (2.3),*

$$\frac{1}{C} |\ln \varepsilon| \leq -a_{ii} \leq C |\ln \varepsilon|, \quad \text{for } n = 3, \quad i = 1, 2.$$

Proof: We consider

$$\bar{w}(x_1, x') = -\frac{x - g(x') - \frac{\varepsilon}{2}}{g(x') - f(x') + \varepsilon} \tag{2.14}$$

on $O_{r/2} := \tilde{\Omega} \cap \{(x_1, x') \mid |x'| < \frac{r}{2}\}$. Use the same proof in Lemma 2.5, we have

$$\int_0^{r/2} \int_{f(x')-\frac{\varepsilon}{2}}^{g(x')+\frac{\varepsilon}{2}} |\partial_{x'} \bar{w}(x_1, x')|^2 dx_1 dx' \leq C.$$

Therefore, it suffices to verify that

$$\int_{\tilde{\Omega} \cap O_{r/2}} |\partial_{x_1} \bar{w}(x_1, x')|^2 \sim |\ln \varepsilon|.$$

Indeed,

$$\begin{aligned} \int_{\tilde{\Omega} \cap O_{r/2}} |\partial_{x_1} \bar{w}(x_1, x')|^2 &= \int_{|x'| < r/2} \frac{1}{g(x') - f(x') + \varepsilon} dx' \\ &\sim \int_0^{r/2} \frac{t}{Ct^2 + \varepsilon} dt \sim |\ln \varepsilon|. \end{aligned}$$

This completes the proof. \square

Lemma 2.7 *Let a_{ii} be defined by (2.3),*

$$\frac{1}{C} \leq -a_{ii} \leq C \quad \text{for } n \geq 4, \quad i = 1, 2.$$

Proof: We only need

$$\int_{O_{r/2}} |\partial_{x_1} \bar{w}(x_1, x')|^2 = \int_{|x'| < r/2} \frac{1}{g(x') - f(x') + \varepsilon} dx' \sim \int_0^{r/2} \frac{t^{n-2}}{Ct^2 + \varepsilon} dt \sim C.$$

The proof is completed. \square

Lemma 2.8 *Let α be defined by (2.6), we have*

$$\frac{1}{C} \leq \alpha \leq C.$$

Proof: By the definition of α and using the second statement in Lemma 2.4, we are done. \square

To summarize, we have

Proposition 2.2 *Let a_{ij} and α be defined by (2.3) and (2.6), we have*

1. $\frac{1}{C\sqrt{\varepsilon}} \leq |a_{11} - \alpha a_{12}| \leq \frac{C}{\sqrt{\varepsilon}} \quad \text{for } n = 2,$
2. $\frac{1}{C} |\ln \varepsilon| \leq |a_{11} - \alpha a_{12}| \leq C |\ln \varepsilon| \quad \text{for } n = 3,$
3. $\frac{1}{C} \leq |a_{11} - \alpha a_{12}| \leq C \quad \text{for } n \geq 4.$

Proof: Since $a_{11} < 0$, $a_{12} > 0$, $a_{11} + a_{12} < 0$ and $\alpha > 0$, we have

$$|a_{11}| < |a_{11} - \alpha a_{12}| < (1 + \alpha)|a_{11}|.$$

Combining the results of Lemma 2.5, Lemma 2.6, Lemma 2.7 and Lemma 2.8, the proof is completed. \square

2.1.3 Estimate of $|b_1 - \alpha b_2|$

Proposition 2.3 *Let b_1 , b_2 , α and $Q_\varepsilon[\varphi]$ be defined by (2.3), (2.6) and (1.10), we have*

$$\frac{|Q_\varepsilon[\varphi]|}{C} \leq |b_1 - \alpha b_2| \leq C\|\varphi\|_{C^2(\partial\Omega)}.$$

Proof: Combining the third result in Lemma 2.4 and Lemma 2.8, we have

$$|b_1 - \alpha b_2| \leq |b_1| + |\alpha||b_2| \leq C\|\varphi\|_{C^2(\partial\Omega)}.$$

On the other hand, by the definition and the harmonicity of v_1 and v_2 and using Lemma 2.4, we obtain

$$\begin{aligned} |b_1 - \alpha b_2| &= \frac{|b_1(a_{22} + a_{12}) - b_2(a_{11} + a_{12})|}{|a_{22} + a_{12}|} \\ &\geq \frac{1}{C} \cdot \left| \int_{\partial D_1} \frac{\partial v_3}{\partial \nu} \int_{\partial\Omega} \frac{\partial v_2}{\partial \nu} - \int_{\partial D_2} \frac{\partial v_3}{\partial \nu} \int_{\partial\Omega} \frac{\partial v_1}{\partial \nu} \right| = \frac{|Q_\varepsilon[\varphi]|}{C}. \end{aligned}$$

This completes the proof. \square

Now we are ready to prove our two main theorems:

Proof of Theorem 1.1-1.2: By Proposition 2.1 and (2.6), then using Proposition 2.2, 2.3, we are done. \square

As we mentioned in Remark 1.1, the strict convexity assumption of the two inclusions can be weakened. In fact, our proofs of Theorem 1.1–1.2 apply, with minor modification, to more general situations:

In \mathbb{R}^n , $n \geq 2$, under the same assumptions in the beginning of Section 1.2 except for the strict convexity condition, $\partial D_1 \cap B(0, r)$ and $\partial D_2 \cap B(0, r)$ can be represented by the graph

of $x_1 = f(x') - \frac{\varepsilon}{2}$ and $x_1 = g(x') + \frac{\varepsilon}{2}$, then $f(0') = g(0') = 0$, $\nabla(g - f)(0') = 0$. Assume further that

$$\lambda_0 |x'|^{2m} \leq g(x') - f(x') \leq \lambda_1 |x'|^{2m}, \quad \forall |x'| \leq r/2, \quad (2.15)$$

for some ε -independent $\lambda_0, \lambda_1 > 0, m \geq 1 \in \mathbb{Z}$.

Under the above assumption, let $u \in H^1(\Omega) \cap C^1(\bar{\Omega})$ be the solution to equation (1.4). For ε sufficiently small, there exist positive constants C and C' , such that

$$\begin{aligned} \frac{|Q_\varepsilon[\varphi]|}{C'} \cdot \varepsilon^{-\frac{n-1}{2m}} &\leq \|\nabla u\|_{L^\infty(\bar{\Omega})} \leq C \|\varphi\|_{C^2(\partial\Omega)} \cdot \varepsilon^{-\frac{n-1}{2m}}, \quad \text{if } n-1 < 2m, \\ \frac{|Q_\varepsilon[\varphi]|}{C'} \cdot \frac{1}{\varepsilon |\ln \varepsilon|} &\leq \|\nabla u\|_{L^\infty(\bar{\Omega})} \leq C \|\varphi\|_{C^2(\partial\Omega)} \cdot \frac{1}{\varepsilon |\ln \varepsilon|}, \quad \text{if } n-1 = 2m, \\ \frac{|Q_\varepsilon[\varphi]|}{C'} \cdot \frac{1}{\varepsilon} &\leq \|\nabla u\|_{L^\infty(\bar{\Omega})} \leq C \|\varphi\|_{C^2(\partial\Omega)} \cdot \frac{1}{\varepsilon}, \quad \text{if } n-1 > 2m, \end{aligned} \quad (2.16)$$

where $Q_\varepsilon[\varphi]$ is defined by (1.10), and C depends on $n, m, \lambda_0, \lambda_1, r_0, \|\partial\Omega\|_{C^{2,\alpha}}, \|\partial D_1\|_{C^{2,\alpha}}$ and $\|\partial D_2\|_{C^{2,\alpha}}$, C' depends on the same as C and also $\|\varphi\|_{C^2(\partial\Omega)}$, but both are independent of ε .

The proof is essentially the same except for the computation of $\int_{\bar{\Omega}} |\nabla v_1|^2$.

In fact,

$$\int_0^{r/2} \int_{f(x')-\frac{\varepsilon}{2}}^{g(x')+\frac{\varepsilon}{2}} |\partial_{x'} \bar{w}(x_1, x')|^2 dx_1 dx' \leq C,$$

still holds. Then by (2.10) and (2.13) we only need to calculate

$$\int_{|x'| < r/2} \frac{1}{g(x') - f(x') + \varepsilon} dx' \sim \int_0^{r/2} \frac{\rho^{n-2}}{\rho^{2m} + \varepsilon} d\rho.$$

Indeed, if $n-1 < 2m$,

$$\int_0^{r/2} \frac{\rho^{n-2}}{\rho^{2m} + \varepsilon} d\rho = \varepsilon^{\frac{n-1}{2m}-1} \int_0^{r/2\varepsilon^{\frac{1}{2m}}} \frac{s^{n-2}}{s^{2m} + 1} ds \sim C \varepsilon^{\frac{n-1}{2m}-1},$$

if $n-1 = 2m$,

$$\int_0^{r/2} \frac{\rho^{n-2}}{\rho^{2m} + \varepsilon} d\rho = \frac{1}{2m} \int_0^{r/2} \frac{1}{\rho^{2m} + \varepsilon} d\rho^{2m} \sim C |\ln \varepsilon|,$$

if $n-1 > 2m$,

$$\int_0^{r/2} \frac{\rho^{n-2}}{\rho^{2m} + \varepsilon} d\rho \sim C.$$

Therefore, we obtain (2.16) by using the same arguments in the proofs of Theorem 1.1 and Theorem 1.2.

Actually, we can replace $2m$ by any real number $\beta > 0$, the results still hold.

2.2 Estimate of $|Q_\varepsilon[\varphi]|$

In order to identify situations when $\|\nabla u\|_{L^\infty}$ behaves exactly as the upper bound established in Theorem 1.1, we estimate in this section $|Q_\varepsilon[\varphi]|$. To emphasize the dependence on ε , we denote D_1, D_2 by $D_{1\varepsilon}, D_{2\varepsilon}$, denote φ by φ_ε , and denote v_1, v_2, v_3 defined by equation (1.7), (1.8), (1.9) as $v_{1\varepsilon}, v_{2\varepsilon}, v_{3\varepsilon}$. In this section we assume, in addition to the hypotheses in Theorem 1.1, that along a sequence $\varepsilon \rightarrow 0$ (we still denote it as ε), $D_{1\varepsilon} \rightarrow D_1^*, D_{2\varepsilon} \rightarrow D_2^*$ in $C^{2,\alpha}$ norm, $\varphi_\varepsilon \rightarrow \varphi^*$ in $C^{1,\alpha}(\partial\Omega)$. We use notation $\tilde{\Omega}^* = \Omega \setminus \overline{D_1^* \cup D_2^*}$, and assume, without loss of generality, that $D_1^* \cap D_2^* = \{0\}$. We will show that as $\varepsilon \rightarrow 0$, $v_{i\varepsilon}$ converges, in appropriate sense, to v_i^* which satisfies

$$\begin{cases} \Delta v_1^* = 0 & \text{in } \tilde{\Omega}^*, \\ v_1^* = 1 & \text{on } \partial D_1^* \setminus \{0\}, \quad v_1^* = 0 & \text{on } \partial\Omega \cup \partial D_2^* \setminus \{0\}, \end{cases} \quad (2.17)$$

$$\begin{cases} \Delta v_2^* = 0 & \text{in } \tilde{\Omega}^*, \\ v_2^* = 1 & \text{on } \partial D_2^* \setminus \{0\}, \quad v_2^* = 0 & \text{on } \partial\Omega \cup \partial D_1^* \setminus \{0\}, \end{cases} \quad (2.18)$$

$$\begin{cases} \Delta v_3^* = 0 & \text{in } \tilde{\Omega}^*, \\ v_3^* = 0 & \text{on } \partial D_1^* \cup \partial D_2^*, \quad v_3^* = \varphi^* & \text{on } \partial\Omega. \end{cases} \quad (2.19)$$

First we prove

Lemma 2.9 *There exist unique $v_i^* \in L^\infty(\tilde{\Omega}^*) \cap C^0(\overline{\tilde{\Omega}^*} \setminus \{0\}) \cap C^2(\tilde{\Omega}^*)$, $i = 1, 2, 3$, which solve equations (2.17), (2.18) and (2.19) respectively. Moreover, $v_i^* \in C^1(\overline{\tilde{\Omega}^*} \setminus \{0\})$.*

Proof: The existence of solutions to the above equations can easily be obtained by Perron's method, see theorem 2.12 and lemma 2.13 in [18]. For reader's convenience, we give below a simple proof of the uniqueness. We only need to prove that 0 is the only solution in $L^\infty(\tilde{\Omega}^*) \cap C^0(\overline{\tilde{\Omega}^*} \setminus \{0\}) \cap C^2(\tilde{\Omega}^*)$ to the following equation:

$$\begin{cases} \Delta w = 0 & \text{in } \tilde{\Omega}^*, \\ w = 0 & \text{on } \partial\tilde{\Omega}^* \setminus \{0\}. \end{cases} \quad (2.20)$$

Indeed, $\forall \varepsilon > 0$, we have

$$|w(x)| \leq \frac{\varepsilon^{n-2} \|w\|_{L^\infty(\tilde{\Omega}^*)}}{|x|^{n-2}}, \quad \text{on } \partial(\tilde{\Omega}^* \setminus B_\varepsilon)(0).$$

By the maximum principle,

$$|w(x)| \leq \frac{\varepsilon^{n-2} \|w\|_{L^\infty(\tilde{\Omega}^*)}}{|x|^{n-2}}, \quad \forall x \in \tilde{\Omega}^* \setminus B_\varepsilon(0).$$

Thus $w \equiv 0$ in $\tilde{\Omega}^*$. The additional regularity $v_i^* \in C^1(\overline{\tilde{\Omega}^*} \setminus \{0\})$ follows from standard elliptic estimates and the regularity of the ∂D_i and $\partial\Omega$. \square

Lemma 2.10 For $i = 1, 2, 3$,

$$v_{i\varepsilon} \longrightarrow v_i^* \quad \text{in } C_{loc}^2(\tilde{\Omega}^*), \quad \text{as } \varepsilon \rightarrow 0, \quad (2.21)$$

$$\int_{\partial\Omega} \frac{\partial v_{i\varepsilon}}{\partial\nu} \longrightarrow \int_{\partial\Omega} \frac{\partial v_i^*}{\partial\nu}, \quad \text{as } \varepsilon \rightarrow 0, \quad i = 1, 2, \quad (2.22)$$

$$\int_{\partial D_{i\varepsilon}} \frac{\partial v_{3\varepsilon}}{\partial\nu} \longrightarrow \int_{\partial D_i^*} \frac{\partial v_3^*}{\partial\nu}, \quad \text{as } \varepsilon \rightarrow 0. \quad (2.23)$$

Proof: By the maximum principle, $\{\|v_{i\varepsilon}\|_{L^\infty}\}$ is bounded by a constant independent of ε . By the uniqueness part of Lemma 2.9, we obtain (2.21) using standard elliptic estimates. By Lemma 2.3, $\{\|\nabla v_{3\varepsilon}\|_{L^\infty}\}$ is bounded by some constant independent of ε , so $\|\nabla v_3^*\|_{L^\infty} < \infty$. Estimate (2.22) and (2.23) follow from standard elliptic estimates. The proof is completed. \square

Similar to $Q_\varepsilon[\varphi_\varepsilon]$, we define

$$Q^*[\varphi^*] := \int_{\partial D_1^*} \frac{\partial v_3^*}{\partial\nu} \int_{\partial\Omega} \frac{\partial v_2^*}{\partial\nu} - \int_{\partial D_2^*} \frac{\partial v_3^*}{\partial\nu} \int_{\partial\Omega} \frac{\partial v_1^*}{\partial\nu}, \quad (2.24)$$

then $Q^* : C^2(\partial\Omega) \mapsto \mathbb{R}$ is a linear functional. Let $Q_\varepsilon[\varphi_\varepsilon]$ and $Q^*[\varphi^*]$ be defined by equation (1.10), (2.24), then, by the above lemmas,

$$Q_\varepsilon[\varphi_\varepsilon] \longrightarrow Q^*[\varphi^*], \quad \text{as } \varepsilon \rightarrow 0.$$

Corollary 2.2 If $\varphi^* \in C^2(\partial\Omega)$ satisfies $Q^*[\varphi^*] \neq 0$, then $|Q_\varepsilon[\varphi_\varepsilon]| \geq \frac{1}{C}$, for some positive constant C which is independent of ε .

In the following we give some examples to show that, in general, the rates of the lower bounds established in Theorem 1.2 are optimal.

Let $\Omega \subset \mathbb{R}^n$, $n \geq 2$, be a bounded open set with $C^{2,\alpha}$ boundary, $0 < \alpha < 1$, which is symmetric with respect to x_1 -variable, i.e., $(x_1, x') \in \Omega$ if and only if $(-x_1, x') \in \Omega$, where $x' = (x_2, \dots, x_n)$.

Let D_1^* be a strictly convex bounded open set in $\{(x_1, x') \in \mathbb{R}^n | x_1 < 0\}$ with $C^{2,\alpha}$ boundary, $0 < \alpha < 1$, satisfying $0 \in \partial D_1^*$ and $\overline{D_1^*} \subset \Omega$. Set $D_2^* = \{(x_1, x') \in \mathbb{R}^n | (-x_1, x') \in D_1^*\}$.

Let $\varphi \in C^2(\partial\Omega) \setminus \{0\}$ satisfy

$$\varphi_{odd}(x_1, x') := \frac{1}{2} [\varphi(x_1, x') - \varphi(-x_1, x')] \leq 0 \text{ (or } \geq 0), \quad (2.25)$$

on $(\partial\Omega)^+ := \{(x_1, x') \in \partial\Omega | x_1 > 0\}$.

For $\varepsilon > 0$ sufficiently small, let

$$\begin{aligned} D_{1\varepsilon} &:= \{(x_1, x') \in \Omega | (x_1 + \frac{\varepsilon}{2}, x') \in D_1^*\}, \\ D_{2\varepsilon} &:= \{(x_1, x') \in \Omega | (x_1 - \frac{\varepsilon}{2}, x') \in D_2^*\}, \\ \varphi_\varepsilon &:= \varphi. \end{aligned}$$

Proposition 2.4 *Under the above assumptions, we have $|Q_\varepsilon[\varphi]| \geq \frac{1}{C}$, for some positive constant C independent of ε . Consequently,*

$$\begin{aligned} \|\nabla u_\varepsilon\|_{L^\infty(\tilde{\Omega})} &\geq \frac{1}{C\sqrt{\varepsilon}} && \text{for } n = 2, \\ \|\nabla u_\varepsilon\|_{L^\infty(\tilde{\Omega})} &\geq \frac{1}{C_\varepsilon |\ln \varepsilon|} && \text{for } n = 3, \\ \|\nabla u_\varepsilon\|_{L^\infty(\tilde{\Omega})} &\geq \frac{1}{C_\varepsilon} && \text{for } n \geq 4, \end{aligned} \quad (2.26)$$

where u_ε is the solution to equation (1.4).

The above proposition can be easily obtained by the following lemma which gives a necessary and sufficient condition instead of condition (2.25) on φ for the lower bounds (2.26) to hold.

Let

$$(v_3^*)_{odd}(x_1, x') := \frac{1}{2} [v_3^*(x_1, x') - v_3^*(-x_1, x')], \quad (2.27)$$

we have

Lemma 2.11 *Under the same hypotheses in Proposition 2.4 except for the condition (2.25), let $Q_\varepsilon[\varphi]$ and $(v_3^*)_{odd}(x)$ be defined by equation (1.10) and (2.27), then the following statements are equivalent:*

1. *For some positive constant C independent of ε , we have $|Q_\varepsilon[\varphi]| \geq \frac{1}{C}$,*
2. *$\int_{\partial D_2^*} \frac{\partial(v_3^*)_{odd}}{\partial\nu} \neq 0$.*

Proof: By symmetry, the strong maximum principle and the Hopf Lemma, we can easily obtain

$$\int_{\partial\Omega} \frac{\partial v_1^*}{\partial\nu} = \int_{\partial\Omega} \frac{\partial v_2^*}{\partial\nu} < 0.$$

Then

$$\begin{aligned} Q^*[\varphi] &= \int_{\partial\Omega} \frac{\partial v_1^*}{\partial\nu} \left(\int_{\partial D_1^*} \frac{\partial v_3^*}{\partial\nu} - \int_{\partial D_2^*} \frac{\partial v_3^*}{\partial\nu} \right) \\ &= \int_{\partial\Omega} \frac{\partial v_1^*}{\partial\nu} \left(\int_{\partial D_1^*} \frac{\partial(v_3^*)_{odd}}{\partial\nu} - \int_{\partial D_2^*} \frac{\partial(v_3^*)_{odd}}{\partial\nu} \right) \\ &= -2 \int_{\partial\Omega} \frac{\partial v_1^*}{\partial\nu} \int_{\partial D_2^*} \frac{\partial(v_3^*)_{odd}}{\partial\nu}. \end{aligned}$$

Hence, $Q^*[\varphi] \neq 0$ if and only if $\int_{\partial D_2^*} \frac{\partial(v_3^*)_{odd}}{\partial\nu} \neq 0$. Then by Corollary 2.2, we complete the proof. \square

Proof of Proposition 2.4: Note that $(v_3^*)_{odd}(0, x') = 0$ by symmetry, and $(v_3^*)_{odd}$ is harmonic with $(v_3^*)_{odd} = \varphi_{odd} \leq 0$ (or ≥ 0) but not identically 0 on $(\partial\Omega)^+$. Now by using the strong maximum principle and the Hopf Lemma, it is clear that $\int_{\partial D_2^*} \frac{\partial(v_3^*)_{odd}}{\partial\nu} \neq 0$. Hence, by Lemma 2.11 and Theorem 1.2, we are done. \square

Remark 2.2 *If $\varphi = \sum_{i=1}^n b_i x_i$ with $b_i \in \mathbb{R}$ and $b_1 \neq 0$, then by Proposition 2.4 we have $|Q_\varepsilon[\varphi]| \geq \frac{1}{C}$. Therefore, by Theorem 1.1 and 1.2, the blow-up rates of $\|\nabla u\|_{L^\infty(\tilde{\Omega})}$ are $\varepsilon^{-1/2}$ in dimension $n = 2$, $(\varepsilon |\ln \varepsilon|)^{-1}$ in dimension $n = 3$ and ε^{-1} in dimension $n \geq 4$.*

Now instead of in a bounded set Ω , we consider in \mathbb{R}^n :

$$\begin{cases} \Delta u_\varepsilon = 0 & \text{in } \mathbb{R}^n \setminus \overline{D_{1\varepsilon} \cup D_{2\varepsilon}}, \\ u_\varepsilon|_+ = u_\varepsilon|_- & \text{on } \partial D_{1\varepsilon} \cup \partial D_{2\varepsilon}, \\ \nabla u_\varepsilon \equiv 0 & \text{in } D_{1\varepsilon} \cup D_{2\varepsilon}, \\ \int_{\partial D_{i\varepsilon}} \frac{\partial u_\varepsilon}{\partial \nu} \Big|_+ = 0 & (i = 1, 2), \\ \limsup_{|x| \rightarrow \infty} |x|^{n-1} |u_\varepsilon(x) - H(x)| < \infty, \end{cases} \quad (2.28)$$

where $H(x)$ is a given entire harmonic function in \mathbb{R}^n .

we have the following result regarding the lower bound for $|\nabla u_\varepsilon|$:

Proposition 2.5 *With the same assumptions on $D_{1\varepsilon}$ and $D_{2\varepsilon}$ as in Proposition 2.4, and let $H(x)$ be an entire harmonic function in \mathbb{R}^n satisfying $H_{\text{odd}}(x_1, x') := \frac{1}{2}[H(x_1, x') - H(-x_1, x')] < 0$ (or > 0) on $\mathbb{R}_+^n := \{(x_1, x') \in \mathbb{R}^n | x_1 > 0\}$, then for some positive constant C independent of ε , we have*

$$\begin{aligned} \|\nabla u_\varepsilon\|_{L^\infty(\mathbb{R}^n \setminus \overline{D_{1\varepsilon} \cup D_{2\varepsilon}})} &\geq \frac{1}{C\sqrt{\varepsilon}} & \text{for } n = 2, \\ \|\nabla u_\varepsilon\|_{L^\infty(\mathbb{R}^n \setminus \overline{D_{1\varepsilon} \cup D_{2\varepsilon}})} &\geq \frac{1}{C\varepsilon |\ln \varepsilon|} & \text{for } n = 3, \\ \|\nabla u_\varepsilon\|_{L^\infty(\mathbb{R}^n \setminus \overline{D_{1\varepsilon} \cup D_{2\varepsilon}})} &\geq \frac{1}{C\varepsilon} & \text{for } n \geq 4, \end{aligned} \quad (2.29)$$

where u_ε is the solution to equation (2.28).

Proof: Step 1. First, we show that there exists a positive constant C independent of ε , such that for any small $\varepsilon > 0$,

$$|x|^{n-1} |u_\varepsilon(x) - H(x)| \leq C, \quad \forall x \in \mathbb{R}^n \setminus \overline{D_{1\varepsilon} \cup D_{2\varepsilon}}. \quad (2.30)$$

(i) For any bounded open set $U \subset \mathbb{R}^n$ with C^1 boundary ∂U satisfying $\partial U \cap \overline{D_{1\varepsilon} \cup D_{2\varepsilon}} = \emptyset$, we have, in view of the first and the fourth lines in (2.28),

$$\int_{\partial U} \frac{\partial u_\varepsilon}{\partial \nu} = \int_{U \setminus \overline{D_{1\varepsilon} \cup D_{2\varepsilon}}} \Delta u_\varepsilon = 0. \quad (2.31)$$

(ii) We show that there exists a positive constant M independent of ε , such that

$$\|u_\varepsilon - H\|_{L^\infty(\mathbb{R}^n \setminus \overline{D_{1\varepsilon} \cup D_{2\varepsilon}})} \leq M, \quad \forall \text{ small } \varepsilon > 0.$$

We only need to prove

$$\|u_\varepsilon - H\|_{L^\infty(\mathbb{R}^n \setminus \overline{D_{1\varepsilon} \cup D_{2\varepsilon}})} \leq \sum_{i=1}^2 (\max_{\overline{D_{i\varepsilon}}} H - \min_{\overline{D_{i\varepsilon}}} H). \quad (2.32)$$

Since $\nabla u_\varepsilon = 0$ in $D_{1\varepsilon} \cup D_{2\varepsilon}$, u_ε is constant on each $D_{i\varepsilon}$, denoted as $C_i(\varepsilon)$. We know that

$$\lim_{|x| \rightarrow \infty} (u_\varepsilon(x) - H(x)) = 0, \quad (2.33)$$

and

$$C_i(\varepsilon) - \max_{\overline{D_{i\varepsilon}}} H \leq u_\varepsilon - H \leq C_i(\varepsilon) - \min_{\overline{D_{i\varepsilon}}} H, \quad \text{on } D_{i\varepsilon}, \quad i = 1, 2. \quad (2.34)$$

If (2.32) did not hold, say,

$$\sup_{\mathbb{R}^n} (u_\varepsilon - H) > \sum_{i=1}^2 (\max_{\overline{D_{i\varepsilon}}} H - \min_{\overline{D_{i\varepsilon}}} H),$$

then, because of (2.33) and (2.34), there would exist $0 < a < \sup_{\mathbb{R}^n} (u_\varepsilon - H)$ such that $U := \{x \in \mathbb{R}^n \mid (u_\varepsilon - H)(x) > a\} \neq \emptyset$ satisfies $\partial U \cap \overline{D_{1\varepsilon} \cup D_{2\varepsilon}} = \emptyset$. We may assume, by the Sard theorem, that a is a regular value of $u_\varepsilon - H$, and therefore ∂U is C^1 . By the Hopf lemma, $\frac{\partial(u_\varepsilon - H)}{\partial \nu} < 0$ on ∂U , and therefore

$$\int_{\partial U} \frac{\partial(u_\varepsilon - H)}{\partial \nu} < 0.$$

On the other hand, using (2.31) and the harmonicity of H in U , we have

$$\int_{\partial U} \frac{\partial(u_\varepsilon - H)}{\partial \nu} = - \int_{\partial U} \frac{\partial H}{\partial \nu} = - \int_U \Delta H = 0.$$

A contradiction.

(iii) Consider $w_\varepsilon(x) := u_\varepsilon(x) - H(x)$. Fix a constant $R_0 > 0$, independent of ε , such that $D_1^* \cup D_2^* \subset B_{R_0/2}(0)$, and let

$$\widetilde{w}_\varepsilon(y) := \frac{1}{|y|^{n-2}} w_\varepsilon\left(\frac{y}{|y|^2}\right), \quad 0 < |y| < \frac{1}{R_0}.$$

Then $\widetilde{w}_\varepsilon$ is harmonic in $B_{1/R_0} \setminus \{0\}$. By the last line of (2.28), there exists a positive constant $C(\varepsilon)$ such that

$$|\widetilde{w}_\varepsilon(y)| \leq C(\varepsilon)|y|, \quad 0 < |y| < \frac{1}{R_0}.$$

Therefore, $\Delta \widetilde{w}_\varepsilon = 0$ in B_{1/R_0} and $\widetilde{w}_\varepsilon(0) = 0$. By (ii), we have $|\widetilde{w}_\varepsilon| \leq C$, on $\partial B_{1/R_0}$, for some positive constant C independent of ε . Hence, $|\widetilde{w}_\varepsilon| \leq C$, $|\nabla \widetilde{w}_\varepsilon| \leq C$ in $B_{1/(2R_0)}$, then

$$|\widetilde{w}_\varepsilon(y)| \leq C|y|, \quad |y| < \frac{1}{2R_0}.$$

Therefore, also using (ii), (2.30) holds.

Step 2. For $R > R_0$, let $\Omega = B_R(0)$. Let $\varphi_\varepsilon := u_\varepsilon|_{\partial\Omega}$, then by Corollary 2.2 and Theorem 1.2 it is enough to show, for some R , that $Q^*[\varphi^*] \neq 0$, where φ^* is defined at the beginning of this section. By symmetry, we have

$$Q^*[\varphi^*] = \int_{\partial\Omega} \frac{\partial v_1^*}{\partial \nu} \left(\int_{\partial D_1^*} \frac{\partial v_3^*}{\partial \nu} - \int_{\partial D_2^*} \frac{\partial v_3^*}{\partial \nu} \right).$$

Without loss of generality, we may assume $H_{\text{odd}}(x) > 0$ on \mathbb{R}_+^n . Recall that v_3^* is the solution of (2.19) with boundary data φ^* . In the following we use notation $(v_3^*)_h$ to denote the the solution of (2.19) with boundary data h . Since $Q^*[\varphi^*]$ is linear on φ^* and by symmetry $Q^*[H_{\text{even}}] = H[\varphi_{\text{even}}^*] = 0$, where $H_{\text{even}}(x) := H(x) - H_{\text{odd}}(x) = \frac{1}{2}[H(x_1, x') + H(-x_1, x')]$ and similar for φ_{even}^* , we may assume $H(x) = H_{\text{odd}}(x)$.

Now consider $w(x) = H(x) - (v_3^*)_H(x)$. Then $w(x)$ is harmonic in $\widetilde{\Omega}^*$ which is defined at the beginning of this section. By symmetry, $w(-x_1, x') = -w(x_1, x')$, $w(x) = H(x)$ on $\partial D_1^* \cup \partial D_2^*$ and $w(x) = 0$ on $\partial\Omega$. Therefore,

$$-2 \int_{\partial D_2^*} H \frac{\partial w}{\partial \nu} = \int_{\widetilde{\Omega}^*} w(x) \Delta w(x) + \int_{\widetilde{\Omega}^*} |\nabla w|^2 = \int_{\widetilde{\Omega}^*} |\nabla w|^2 \geq 0.$$

On the other hand, $(v_3^*)_H = 0$ on ∂D_2^* , $(v_3^*)_H > 0$ on $(\partial\Omega)^+$ and, by the oddness of $(v_3^*)_H$, $(v_3^*)_H = 0$ on $\{(x_1, x') \mid x_1 = 0\}$. Thus, by the maximum principle and the strong maximum principle, $(v_3^*)_H > 0$ in $\widetilde{\Omega}^*$ and in turn, using the Hopf lemma, $\frac{\partial (v_3^*)_H}{\partial \nu} > 0$ on ∂D_2^* . Hence, using the harmonicity of H ,

$$\begin{aligned} \max_{\partial D_2^*} H \int_{\partial D_2^*} \frac{\partial (v_3^*)_H}{\partial \nu} &\geq \int_{\partial D_2^*} H \frac{\partial (v_3^*)_H}{\partial \nu} \geq \int_{\partial D_2^*} H \frac{\partial H}{\partial \nu} - \int_{\partial D_2^*} H \frac{\partial w}{\partial \nu} \\ &\geq \int_{D_2^*} |\nabla H|^2 \geq \frac{1}{C}, \end{aligned}$$

Therefore,

$$\int_{\partial D_2^*} \frac{\partial (v_3^*)_H}{\partial \nu} \geq \frac{1}{C},$$

for positive constant C independent of R .

For $s_\varepsilon := \varphi_\varepsilon - H$ on $\partial\Omega$, by step 1, there exists a constant $C > 0$ which is independent of ε and R , such that $\|s_\varepsilon\|_{L^\infty(\partial\Omega)} \leq CR^{1-n}$. By Remark 2.1, we have $\|\nabla(v_3^*)_{s^*}\|_{L^\infty(\partial D_1^* \cup \partial D_2^*)} \leq C\|s^*\|_{L^\infty(\partial\Omega)}$, thus,

$$\left| \int_{\partial D_i^*} \frac{\partial(v_3^*)_{s^*}}{\partial\nu} \right| \leq C \int_{\partial D_i^*} \|s^*\|_{L^\infty(\partial\Omega)} \leq CR^{1-n},$$

for some positive constant C independent of ε and R .

Therefore, for large enough R ,

$$\int_{\partial D_2^*} \frac{\partial(v_3^*)_{\varphi^*}}{\partial\nu} = \int_{\partial D_2^*} \frac{\partial(v_3^*)_H}{\partial\nu} + \int_{\partial D_2^*} \frac{\partial(v_3^*)_{s^*}}{\partial\nu} \geq \frac{1}{C} \neq 0.$$

It is also clear that $\int_{\partial\Omega} \frac{\partial v_1^*}{\partial\nu} < 0$. Thus,

$$Q^*[\varphi^*] = -2 \int_{\partial\Omega} \frac{\partial v_1^*}{\partial\nu} \int_{\partial D_2^*} \frac{\partial(v_3^*)_{\varphi^*}}{\partial\nu} \neq 0.$$

This proof is completed. □

Remark 2.3 In \mathbb{R}^2 , when $D_{1\varepsilon}$ and $D_{2\varepsilon}$ are identical balls of radius 1, the estimate (2.29) was established in [6] under a weaker assumption $\partial_{x_1}H(0) \neq 0$.

2.3 Proof of Theorem 1.3 and 1.4

In the introduction, similar to the harmonic case, we still decompose $u = C_1V_1 + C_2V_2 + V_3$ as in (1.14).

Proposition 2.1 holds since Lemma 2.1–2.3 hold for V_1, V_2, V_3 defined by (1.15)–(1.17) and $\rho \in C^2(\tilde{\Omega})$ which is the solution to:

$$\begin{cases} \partial_{x_j} \left(a_2^{ij}(x) \partial_{x_i} \rho \right) = 0 & \text{in } \tilde{\Omega}, \\ \rho = 0 \text{ on } \partial D_1 \cup \partial D_2, \quad \rho = 1 \text{ on } \partial\Omega. \end{cases}$$

The proofs are essentially the same.

Now we start to estimate $|C_1 - C_2|$. By the decomposition formula (1.14), instead of (2.3), we denote

$$\begin{aligned} a_{lm} &= \int_{\partial D_l} a_2^{ij}(x) \partial_{x_i} V_m \nu_j \quad (l, m = 1, 2), \\ b_l &= \int_{\partial D_l} a_2^{ij}(x) \partial_{x_i} V_3 \nu_j \quad (l = 1, 2). \end{aligned} \quad (2.35)$$

Then Lemma 2.4 and (2.4)–(2.6) still hold for a_{lm} and b_l defined above.

In fact, to prove Lemma 2.4 with general coefficients, we only need to change $\frac{\partial^*}{\partial \nu}$ to $a_2^{ij}(x) \partial_{x_i}^* \nu_j$, change Δ^* in $\partial_{x_j} \left(a_2^{ij}(x) \partial_{x_i}^* \right)$ and change v_1, v_2, v_3 in V_1, V_2, V_3 , respectively, in the original proof of Lemma 2.4. For instance, (2.7) is changed to

$$\begin{aligned} 0 &= \int_{\bar{\Omega}} \partial_{x_j} \left(a_2^{ij}(x) \partial_{x_i} V_1 \right) \cdot V_2 - \int_{\bar{\Omega}} \partial_{x_j} \left(a_2^{ij}(x) \partial_{x_i} V_2 \right) \cdot V_1 \\ &= - \int_{\partial D_2} a_2^{ij}(x) \partial_{x_i} V_1 \nu_j \cdot 1 + \int_{\partial D_1} a_2^{ij}(x) \partial_{x_i} V_2 \nu_j \cdot 1 \\ &= -a_{21} + a_{12}. \end{aligned} \quad (2.36)$$

Therefore, to estimate $|C_1 - C_2|$, it is equivalent to estimating $|a_{11} - \alpha a_{12}|$ and $|b_1 - \alpha b_2|$.

For $|a_{11} - \alpha a_{12}|$, Lemma 2.5–2.7 still hold for $a_{ll}(l = 1, 2)$ defined by (2.35). The proof is quite similar and the only thing which needs to be shown is the following:

$$\begin{aligned} 0 &= \int_{\bar{\Omega}} \partial_{x_j} \left(a_2^{ij}(x) \partial_{x_i} V_1 \right) \cdot V_1 \\ &= - \int_{\bar{\Omega}} a_2^{ij}(x) \partial_{x_i} V_1 \partial_{x_j} V_1 - \int_{\partial D_1} a_2^{ij}(x) \partial_{x_i} V_1 \nu_j \cdot 1 \\ &= - \int_{\bar{\Omega}} a_2^{ij}(x) \partial_{x_i} V_1 \partial_{x_j} V_1 - a_{11}, \end{aligned}$$

i.e.

$$a_{11} = - \int_{\bar{\Omega}} a_2^{ij}(x) \partial_{x_i} V_1 \partial_{x_j} V_1.$$

Then by the uniform ellipticity of $a_2^{ij}(x)$ and the harmonicity of v_1 ,

$$|a_{11}| \geq \lambda \int_{\bar{\Omega}} |\nabla V_1|^2 \geq \lambda \int_{\bar{\Omega}} |\nabla v_1|^2,$$

and

$$|a_{11}| \leq \int_{\bar{\Omega}} a_2^{ij}(x) \partial_{x_i} w \partial_{x_j} w \leq \Lambda \int_{\bar{\Omega}} |\nabla w|^2 \leq \Lambda \int_{\bar{\Omega} \cap O_{\tau/2}} |\nabla \bar{w}|^2 + C,$$

where w is defined in the proof of Lemma 2.5 with the same boundary data of V_1 and \overline{w} is defined by (2.9) and (2.14).

Thus, Lemma 2.5–2.7 follow by the same computations. Then Lemma 2.8 and Proposition 2.2 hold with the same proofs.

For $|b_1 - \alpha b_2|$, Proposition 2.3 also holds for $b_l (l = 1, 2)$ defined by (2.35) and $Q_\varepsilon[\varphi]$ defined by (1.18). The proof is the same after changing $\frac{\partial^*}{\partial \nu}$ to $a_2^{ij}(x) \partial_{x_i}^* \nu_j$.

Combining the above propositions, we obtain our theorems.

2.4 Appendix A: Some elementary results for the perfect conductivity problem

Assume that in \mathbb{R}^n , Ω and ω are bounded open sets with $C^{2,\alpha}$ boundaries, $0 < \alpha < 1$, satisfying

$$\overline{\omega} = \bigcup_{s=1}^m \overline{\omega}_s \subset \Omega,$$

where $\{\omega_s\}$ are connected components of ω . Clearly, $m < \infty$ and ω_s is open for all $1 \leq s \leq m$. Given $\varphi \in C^2(\partial\Omega)$, the conductivity problem we consider is the following transmission problem with Dirichlet boundary condition:

$$\begin{cases} \partial_{x_j} \left\{ \left[(k a_1^{ij}(x) - a_2^{ij}(x)) \chi_\omega + a_2^{ij}(x) \right] \partial_{x_i} u_k \right\} = 0 & \text{in } \Omega, \\ u_k = \varphi & \text{on } \partial\Omega, \end{cases} \quad (2.37)$$

where $k = 1, 2, 3, \dots$, and χ_ω is the characteristic function of ω .

The $n \times n$ matrixes $A_1(x) := (a_1^{ij}(x))$ in ω , $A_2(x) := (a_2^{ij}(x))$ in $\Omega \setminus \overline{\omega}$ are symmetric and \exists a constant $\Lambda \geq \lambda > 0$ such that

$$\lambda |\xi|^2 \leq a_1^{ij}(x) \xi_i \xi_j \leq \Lambda |\xi|^2 \quad (\forall x \in \omega), \quad \lambda |\xi|^2 \leq a_2^{ij}(x) \xi_i \xi_j \leq \Lambda |\xi|^2 \quad (\forall x \in \Omega \setminus \omega)$$

for all $\xi \in \mathbb{R}^n$ and $a_1^{ij}(x) \in C^2(\overline{\omega})$, $a_2^{ij}(x) \in C^2(\overline{\Omega} \setminus \omega)$.

Equation (2.37) can be rewritten in the following form to emphasize the transmission condition on $\partial\omega$:

$$\left\{ \begin{array}{ll} \partial_{x_j} \left(a_1^{ij}(x) \partial_{x_i} u_k \right) = 0 & \text{in } \omega, \\ \partial_{x_j} \left(a_2^{ij}(x) \partial_{x_i} u_k \right) = 0 & \text{in } \Omega \setminus \overline{\omega}, \\ u_k|_+ = u_k|_-, & \text{on } \partial\omega, \\ a_2^{ij}(x) \partial_{x_i} u_k \nu_j|_+ = k a_1^{ij}(x) \partial_{x_i} u_k \nu_j|_- & \text{on } \partial\omega, \\ u_k = \varphi & \text{on } \partial\Omega. \end{array} \right. \quad (2.38)$$

We list the following results which are well known and omit the proofs.

Theorem 2.1 *If $u_k \in H^1(\Omega)$ is a solution of equation (2.37), then $u_k \in C^1(\overline{\Omega \setminus \omega}) \cap C^1(\overline{\omega})$ and satisfies equation (2.38).*

If $u_k \in C^1(\overline{\Omega \setminus \omega}) \cap C^1(\overline{\omega})$ is a solution of equation (2.38), then $u_k \in H^1(\Omega)$ and satisfies equation (2.37).

Theorem 2.2 *There exists at most one solution $u_k \in H^1(\Omega)$ to equation (2.37).*

The existence of the solution can be obtained by using the variational method. For every k , we define the energy functional

$$I_k[v] := \frac{k}{2} \int_{\omega} a_1^{ij}(x) \partial_{x_i} v \partial_{x_j} v + \frac{1}{2} \int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} v \partial_{x_j} v, \quad (2.39)$$

where v belongs to the set

$$H_{\varphi}^1(\Omega) := \{v \in H^1(\Omega) \mid v = \varphi \text{ on } \partial\Omega\}.$$

Theorem 2.3 *For every k , there exists a minimizer $u_k \in H^1(\Omega)$ satisfying*

$$I_k[u_k] = \min_{v \in H_{\varphi}^1(\Omega)} I_k[v].$$

Moreover, $u_k \in H^1(\Omega)$ is a solution of equation (2.37).

Comparing equation (2.38), when $k = +\infty$, the perfectly conducting problem turns out

to be:

$$\begin{cases} \partial_{x_j} \left(a_2^{ij}(x) \partial_{x_i} u \right) = 0 & \text{in } \Omega \setminus \overline{\omega}, \\ u|_+ = u|_- & \text{on } \partial\omega, \\ \nabla u = 0 & \text{in } \omega, \\ \int_{\partial\omega_s} a_2^{ij}(x) \partial_{x_i} u \nu_j|_+ = 0 & (s = 1, 2, \dots, m), \\ u = \varphi & \text{on } \partial\Omega. \end{cases} \quad (2.40)$$

We also have similar results:

Theorem 2.4 *If $u \in H^1(\Omega)$ satisfies equation (2.40) except for the fourth line, then $u \in C^1(\overline{\Omega \setminus \omega}) \cap C^1(\overline{\omega})$.*

Proof: By the third line of equation (2.40), we have $u \equiv \text{const}$ on each component of ω , so $u \equiv \text{const}$ on each component of $\partial\omega$. Thus $u \equiv \text{const}$ on each component of $\partial(\Omega \setminus \overline{\omega})$.

Since $u \in H^1(\Omega)$ satisfies $\partial_{x_i} \left(a_2^{ij}(x) \partial_{x_i} u \right) = 0$ in $\Omega \setminus \overline{\omega}$, $u|_{\partial\Omega} = \varphi \in C^2(\partial\Omega)$ and $u \equiv \text{const}$ on each component of $\partial(\Omega \setminus \overline{\omega})$, by the elliptic regularity theory, we have $u \in C^1(\overline{\Omega \setminus \omega}) \cap C^1(\overline{\omega})$. \square

Theorem 2.5 *There exists at most one solution $u \in H^1(\Omega) \cap C^1(\overline{\Omega \setminus \omega}) \cap C^1(\overline{\omega})$ of equation (2.40).*

Proof: It is equivalent to showing that if $\varphi = 0$, equation (2.40) only has the solution $u \equiv 0$.

Integrating by parts in the first line of equation (2.40), we have

$$\begin{aligned} 0 &= - \int_{\Omega \setminus \overline{\omega}} \partial_{x_j} \left(a_2^{ij}(x) \partial_{x_i} u \right) \cdot u \\ &= \int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} u \partial_{x_j} u - \int_{\partial\Omega} u \cdot a_2^{ij}(x) \partial_{x_i} u \nu_j|_- + \int_{\partial\omega} u \cdot a_2^{ij}(x) \partial_{x_i} u \nu_j|_+ \\ &\geq \lambda \int_{\Omega \setminus \overline{\omega}} |\nabla u|^2 - \int_{\partial\Omega} \varphi \cdot a_2^{ij}(x) \partial_{x_i} u \nu_j|_- + C_s \int_{\partial\omega_s} a_2^{ij}(x) \partial_{x_i} u \nu_j|_+ \\ &= \lambda \int_{\Omega \setminus \overline{\omega}} |\nabla u|^2. \end{aligned}$$

Thus $\nabla u = 0$ in $\Omega \setminus \overline{\omega}$. And since $u = \varphi = 0$ on $\partial\Omega$, we have $u \equiv 0$ in $\Omega \setminus \overline{\omega}$. Since $u|_+ = u|_-$ on $\partial\omega$ and $u \equiv C$ on $\overline{\omega}$, we get $u = 0$ on $\overline{\omega}$. Hence $u \equiv 0$ in Ω , i.e. $u \equiv 0$ is the only solution of (2.40) when $\varphi = 0$. \square

Define the energy functional

$$I_\infty[v] := \frac{1}{2} \int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} v \partial_{x_j} v, \quad (2.41)$$

where v belongs to the set

$$\mathcal{A} := \{v \in H_\varphi^1(\Omega) \mid \nabla v \equiv 0 \text{ in } \omega\}.$$

Theorem 2.6 *There exists a minimizer $u \in \mathcal{A}$ satisfying*

$$I_\infty[u] = \min_{v \in \mathcal{A}} I_\infty[v].$$

Moreover, $u \in H^1(\Omega) \cap C^1(\overline{\Omega \setminus \omega}) \cap C^1(\overline{\omega})$ is a solution of equation (2.40).

Proof: By the lower-semi continuity of I_∞ and the weakly closed property of \mathcal{A} , it is easy to see that the minimizer $u \in \mathcal{A}$ exists and satisfies $\partial_{x_j} (a_2^{ij}(x) \partial_{x_i} u) = 0$ in $\Omega \setminus \overline{\omega}$. The only thing which needs to be shown is the fourth line in equation (2.40), i.e.

$$\int_{\partial \omega_s} a_2^{ij}(x) \partial_{x_i} u \nu_j|_+ = 0, \quad s = 1, 2, \dots, m.$$

In fact, since u is a minimizer, for any $\phi \in C_c^\infty(\Omega)$ satisfying $\phi \equiv 1$ on $\overline{\omega}_s$ and $\phi \equiv 0$ on $\overline{\omega}_t (t \neq s)$, let

$$i(t) := I_\infty[u + t\phi] \quad (t \in \mathbb{R}),$$

we have

$$i'(0) := \left. \frac{di}{dt} \right|_{t=0} = \int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} u \phi_{x_j} = 0.$$

Therefore

$$\begin{aligned} 0 &= - \int_{\Omega \setminus \overline{\omega}} \partial_{x_j} (a_2^{ij}(x) \partial_{x_i} u_k) \phi \\ &= \int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} u \phi_{x_j} + \int_{\partial \omega_s} \phi \cdot a_2^{ij}(x) \partial_{x_i} u \nu_j|_+ \\ &= \int_{\partial \omega_s} a_2^{ij}(x) \partial_{x_i} u \nu_j|_+, \end{aligned}$$

for all $s = 1, 2, \dots, m$. □

Finally, we give the relationship between u_k and u .

Theorem 2.7 *Let u_k and u in $H^1(\Omega)$ be the solutions of equations (2.38) and (2.40), respectively. Then*

$$u_k \rightharpoonup u \text{ in } H^1(\Omega), \quad \text{as } k \rightarrow +\infty,$$

and

$$\lim_{k \rightarrow +\infty} I_k[u_k] = I_\infty[u],$$

where I_k and I_∞ are defined as (2.39) and (2.41).

Proof: Step 1. By the uniqueness of the solution to equation (2.40), we only need to show that there exists a weak limit u of a subsequence of $\{u_k\}$ in $H^1(\Omega)$ and u is the solution of equation (2.40).

(1) To show that after passing to a subsequence, u_k weakly converges in $H^1(\Omega)$ to some u .

Let $\eta \in H_\varphi^1(\Omega)$ be fixed and satisfy $\eta \equiv 0$ on $\bar{\omega}$, then since u_k is the minimizer of I_k in $H_\varphi^1(\Omega)$, we have

$$\frac{\lambda}{2} \|\nabla u_k\|_{L^2(\Omega)}^2 \leq I_k[u_k] \leq I_k[\eta] = \frac{1}{2} \int_{\Omega \setminus \bar{\omega}} a_2^{ij}(x) \eta_{x_i} \eta_{x_j} \leq \frac{\Lambda}{2} \|\eta\|_{H^1(\Omega)}^2,$$

i.e.

$$\|\nabla u_k\|_{L^2(\Omega)} \leq \|\eta\|_{H^1(\Omega)} \doteq \overline{M},$$

where \overline{M} is independent of k .

Since $u_k = \varphi$ on $\partial\Omega$ and $\sup_k \|u_k\|_{H^1(\Omega)} < \infty$, we have $u_k \rightharpoonup u$ in $H_\varphi^1(\Omega)$.

(2) To show that u is a solution of equation (2.40).

In fact, we only need to prove the following three conditions:

$$\partial_{x_j} \left(a_2^{ij}(x) \partial_{x_i} u \right) = 0 \quad \text{in } \Omega \setminus \bar{\omega}, \quad (2.42)$$

$$\nabla u = 0 \quad \text{in } \omega, \quad (2.43)$$

$$\int_{\partial\omega_s} a_2^{ij}(x) \partial_{x_i} u_k \nu_j|_+ = 0, \quad s = 1, 2, \dots, m. \quad (2.44)$$

(i) For every k , since $u_k \in H^1(\Omega)$ is the solution of equation (2.37), then

$\forall \phi \in C_c^\infty(\Omega)$, we have

$$k \int_{\omega} a_1^{ij}(x) \partial_{x_i} u_k \phi_{x_j} + \int_{\Omega \setminus \bar{\omega}} a_2^{ij}(x) \partial_{x_i} u_k \phi_{x_j} = 0.$$

Thus, $\forall \phi \in C_c^\infty(\Omega \setminus \overline{\omega}) \subset C_c^\infty(\Omega)$,

$$0 = \int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} u_k \phi_{x_j} \longrightarrow \int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} u \phi_{x_j},$$

since $u_k \rightharpoonup u$ in $H_\varphi^1(\Omega) \subset H^1(\Omega)$.

Therefore,

$$\int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} u \phi_{x_j} = 0, \quad \forall \phi \in C_c^\infty(\Omega \setminus \overline{\omega}),$$

i.e. (2.42).

(ii) Let $\eta \in H_\varphi^1(\Omega)$ be fixed and satisfy $\eta \equiv 0$ on $\overline{\omega}$, then since u_k is the minimizer of I_k in $H_\varphi^1(\Omega)$, we have

$$\frac{k\lambda}{2} \|\nabla u_k\|_{L^2(\omega)}^2 \leq I_k[u_k] \leq I_k[\eta] = \frac{1}{2} \int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} \eta \partial_{x_j} \eta \leq \frac{\Lambda}{2} \|\eta\|_{H^1(\Omega)}^2,$$

which implies

$$\|\nabla u_k\|_{L^2(\omega)}^2 \rightarrow 0, \quad \text{as } k \rightarrow \infty.$$

By (1), since $u_k \rightharpoonup u$ in $H^1(\Omega)$, then $u_k \rightharpoonup u$ in $H^1(\omega)$. Therefore, by the lower-semi continuity, we get

$$\begin{aligned} 0 &\leq \lambda \int_{\omega} |\nabla u|^2 \leq \int_{\omega} a_1^{ij}(x) \partial_{x_i} u \partial_{x_j} u \leq \int_{\omega} a_1^{ij}(x) \partial_{x_i} u_k \partial_{x_j} u_k \\ &\leq \Lambda \|\nabla u_k\|_{L^2(\omega)}^2 \longrightarrow 0, \quad \text{as } k \longrightarrow \infty. \end{aligned}$$

Hence, $\int_{\omega} |\nabla u|^2 = 0 \implies \nabla u \equiv 0$ in ω , which is just (2.43).

(iii) By (i) and (ii), u satisfies (2.42) and is either constant or φ on each component of $\partial(\Omega \setminus \overline{\omega})$. Thus, $u \in C^2(\overline{\Omega \setminus \omega})$. For each $s = 1, 2, \dots, m$, we construct a function $\varrho \in C^2(\overline{\Omega \setminus \omega})$, such that $\varrho = 1$ on $\partial\omega_s$, $\varrho = 0$ on $\partial\omega_t$ ($t \neq s$), and $\varrho = 0$ on $\partial\Omega$.

By Green's Identity, we have the following:

$$\begin{aligned} 0 &= - \int_{\Omega \setminus \overline{\omega}} \partial_{x_j} \left(a_2^{ij}(x) \partial_{x_i} u_k \right) \varrho \\ &= \int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} u_k \partial_{x_j} \varrho - \int_{\partial\Omega} \varrho a_2^{ij}(x) \partial_{x_i} u_k \nu_j|_- + \int_{\partial\omega} \varrho a_2^{ij}(x) \partial_{x_i} u_k \nu_j|_+ \\ &= \int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} u_k \partial_{x_j} \varrho + k \int_{\partial\omega_s} a_1^{ij}(x) \partial_{x_i} u_k \nu_j|_- \\ &= \int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} u_k \partial_{x_j} \varrho. \end{aligned}$$

Similarly,

$$0 = - \int_{\Omega \setminus \overline{\omega}} \partial_{x_j} \left(a_2^{ij}(x) \partial_{x_i} u \right) \varrho = \int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} u \partial_{x_j} \varrho + \int_{\partial \omega_s} a_2^{ij}(x) \partial_{x_i} u \nu_j|_+.$$

Since $u_k \rightharpoonup u$ in $H^1(\Omega)$, it follows

$$0 = \int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} u_k \partial_{x_j} \varrho \longrightarrow \int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} u \partial_{x_j} \varrho.$$

Thus,

$$\int_{\partial \omega_s} a_2^{ij}(x) \partial_{x_i} u \nu_j|_+ = 0,$$

for any $s = 1, 2, \dots, m$. Therefore, we finish the proof of the first part.

Step 2. Since u_k is a minimizer of I_k and $\nabla u = 0$ in ω , for any $k \in \mathbb{N}$,

$$I_k[u_k] \leq I_k[u] = I_\infty[u].$$

Then $\limsup_{k \rightarrow +\infty} I_k[u_k] \leq I_\infty[u]$.

On the other hand, by Theorem 2.7, since u is the weak limit of $\{u_k\}$ in $H^1(\Omega)$, we obtain

$$I_\infty[u] = \int_{\Omega} a_2^{ij}(x) \partial_{x_i} u \partial_{x_j} u \leq \liminf_{k \rightarrow +\infty} \int_{\Omega} a_2^{ij}(x) \partial_{x_i} u_k \partial_{x_j} u_k \leq \liminf_{k \rightarrow +\infty} I_k[u_k].$$

Therefore,

$$\lim_{k \rightarrow +\infty} I_k[u_k] = I_\infty[u].$$

The proof is completed. □

Chapter 3

The perfect and insulated conductivity problems with multiple inclusions

This chapter studies the gradient estimates for the perfect and insulated conductivity problems with multiple inclusions as stated in the introduction. We will first prove Theorem 1.5–1.6 in the case, in equation (1.21) and (1.24), $a_2^{ij}(x) \equiv \delta_{ij}$ where $\delta_{ij} = 0$ if $i \neq j$ and $\delta_{ij} = 1$ if $i = j$.

3.1 The perfect conductivity problem with multiple inclusions

Instead of considering the equation with general coefficients as stated in the introduction, we first prove Theorem 1.5 for the following perfect conductivity problem:

$$\left\{ \begin{array}{ll} \Delta u = 0 & \text{in } \tilde{\Omega} := \Omega \setminus \overline{\bigcup_{i=1}^m D_i}, \\ u|_+ = u|_- & \text{on } \partial D_i, \ (i = 1, 2, \dots, m), \\ \nabla u \equiv 0 & \text{in } D_i \ (i = 1, 2, \dots, m), \\ \int_{\partial D_i} \frac{\partial u}{\partial \nu} \Big|_+ = 0 & (i = 1, 2, \dots, m), \\ u = \varphi & \text{on } \partial \Omega. \end{array} \right. \quad (3.1)$$

The existence and uniqueness of solutions to equation (3.1) are well known, for reference, see Appendix A in chapter 2. In this part we establish the gradient estimates for solutions to equation (3.1). Comparing with the results in chapter 2, where we gave gradient estimates of u in the case of two inclusions, here we manage to establish more accurate local gradient estimates of u in the case of multiple inclusions, as stated in Theorem 1.5.

Note that the gradient of u only blows up in the small area between D_i and D_j when the distance ε_{ij} between D_i and D_j approaches to 0. As a consequence of Theorem 1.5, we have the global estimates of u in $\tilde{\Omega}$.

Corollary 3.1 *Let $\Omega, \{D_i\} \subset \mathbb{R}^n$, $\{\varepsilon_{ij}\}$ be defined as in (1.19), $\varepsilon := \min_{i \neq j} \varepsilon_{ij} > 0$, and $\varphi \in C^2(\partial\Omega)$. Let $u \in H^1(\Omega) \cap C^1(\bar{\Omega})$ be the solution to equation (3.1). For ε sufficiently small, there is a positive constant C which depends only on $n, \kappa_0, r_0, \|\partial\Omega\|_{C^{2,\alpha}}, \{\|\partial D_i\|_{C^{2,\alpha}}\}$, but independent of ε such that*

$$\begin{aligned} \|\nabla u\|_{L^\infty(\bar{\Omega})} &\leq \frac{C}{\sqrt{\varepsilon}} \|\varphi\|_{C^2(\partial\Omega)} && \text{for } n = 2, \\ \|\nabla u\|_{L^\infty(\bar{\Omega})} &\leq \frac{C}{\varepsilon |\ln \varepsilon|} \|\varphi\|_{C^2(\partial\Omega)} && \text{for } n = 3, \\ \|\nabla u\|_{L^\infty(\bar{\Omega})} &\leq \frac{C}{\varepsilon} \|\varphi\|_{C^2(\partial\Omega)} && \text{for } n \geq 4. \end{aligned} \tag{3.2}$$

Note that from equation (3.1), we know that $u \equiv C_i$ on ∂D_i , $1 \leq i \leq m$, where $\{C_i\}$ are some unknown constants. In order to prove Theorem 1.5, first we establish the estimates of $|C_i - C_j|$ for any $1 \leq i \neq j \leq m$, which later will allow us to control the gradient of u in the small area between D_i and D_j .

3.1.1 A Matrix Theorem

In the process of estimating $|C_i - C_j|$, the following theorem plays a crucial role.

Let $s \geq 1$ be an integer, and let $P \in \mathbb{R}^{s \times s}$, $\alpha, \beta \in \mathbb{R}^s$,

$$P = \begin{pmatrix} p_{11} & p_{12} & \cdots & p_{1s} \\ p_{21} & p_{22} & \cdots & p_{2s} \\ \vdots & \vdots & \ddots & \vdots \\ p_{s1} & p_{s2} & \cdots & p_{ss} \end{pmatrix}, \quad \alpha = \begin{pmatrix} \alpha_1 \\ \alpha_2 \\ \vdots \\ \alpha_s \end{pmatrix}, \quad \beta = \begin{pmatrix} \beta_1 \\ \beta_2 \\ \vdots \\ \beta_s \end{pmatrix}, \tag{3.3}$$

and satisfy, for $i, j = 1, 2, \dots, s$

$$\begin{aligned} (1) \quad & p_{ii} > 0, \quad p_{ij} = p_{ji} < 0 \quad (i \neq j), \quad \frac{1}{C_0} \leq \bar{p}_i := \sum_{j=1}^s p_{ij} \leq C_0, \\ (2) \quad & |\beta_i| \leq C_0, \end{aligned} \tag{3.4}$$

where C_0 is a universal positive constant.

Then we have

Theorem 3.1 For an integer $s \geq 1$, let $P \in \mathbb{R}^{s \times s}$, $\alpha, \beta \in \mathbb{R}^s$ be defined as in (3.3) and satisfy (3.4). Let α be the solution of the following system of linear equations

$$P\alpha = \beta. \quad (3.5)$$

Then there exists a constant $C := C(C_0) > 0$ such that

$$|\alpha_i - \alpha_j| \leq \frac{C}{|p_{ij}|}. \quad (3.6)$$

Before proving the above theorem we first give the following lemmas.

Lemma 3.1 For an integer $t \geq 2$, let $Q_{t \times (t-1)} \in \mathbb{R}^{t \times (t-1)}$ be a matrix as follows

$$\begin{pmatrix} q_{12} & q_{13} & \cdots & q_{1t} \\ q_{22} & q_{23} & \cdots & q_{2t} \\ q_{32} & q_{33} & \cdots & q_{3t} \\ \vdots & \vdots & \ddots & \vdots \\ q_{t2} & q_{t3} & \cdots & q_{tt} \end{pmatrix} \quad (3.7)$$

and it satisfies for $i = 1, 2, \dots, t$ and $j = 2, 3, \dots, t$

$$q_{jj} > 0, \quad q_{ij} = q_{ji} < 0 \quad (i \neq j) \quad \text{and} \quad \bar{q}_j := \sum_{i=1}^t q_{ij} > 0. \quad (3.8)$$

Let $Q_i \in \mathbb{R}^{(t-1) \times (t-1)}$ ($i = 1, 2, \dots, t$) denote the matrix obtained from eliminating the i^{th} row of the above matrix $Q_{t \times (t-1)}$. Then the sign of $\det Q_i$ is $(-1)^{i-1}$ and

$$\det Q_1 = \max_{i=1,2,\dots,t} |\det Q_i|.$$

Proof: Step 1. We first show that the sign of $\det Q_i$ is $(-1)^{i-1}$ for $1 \leq i \leq t$.

For $i = 1$, Q_1 is a $(t-1) \times (t-1)$ diagonal dominant matrix. Since each diagonal element is positive we know that the sign of $\det Q_1$ is positive.

Next, we use induction by t to show that for any $t \geq 2$, $\det Q_i$ is $(-1)^{i-1}$ for $2 \leq i \leq t$.

1) For $t = 2$, $\det Q_2 = q_{12} < 0$.

2) Assume that for $t = l - 1$, the sign of $\det Q_i$ is $(-1)^{i-1}$ for any $2 \leq i \leq l - 1$.

For $t = l$,

$$\begin{aligned}
\det Q_2 &= \begin{vmatrix} q_{12} & q_{13} & \cdots & q_{1t} \\ q_{32} & q_{33} & \cdots & q_{3t} \\ \vdots & \vdots & \ddots & \vdots \\ q_{t2} & q_{t3} & \cdots & q_{tt} \end{vmatrix} \\
&= q_{12} \det Q_{12} + (-1)q_{32} \det Q_{22} + \cdots + (-1)^{l-2} q_{l2} \det Q_{(l-1)2}, \tag{3.9}
\end{aligned}$$

where $Q_{i2} \in \mathbb{R}^{(l-2) \times (l-2)}$ ($i = 1, 2, \dots, l-1$) is obtained from eliminating the i^{th} row of the following $(l-1) \times (l-2)$ matrix

$$\begin{pmatrix} q_{13} & q_{14} & \cdots & q_{1l} \\ q_{33} & q_{34} & \cdots & q_{3l} \\ q_{43} & q_{44} & \cdots & q_{4l} \\ \vdots & \vdots & \ddots & \vdots \\ q_{l3} & q_{l4} & \cdots & q_{ll} \end{pmatrix}.$$

By induction the sign of $\det Q_{i2}$ is $(-1)^{i-1}$. Thus the i^{th} term of (3.9) has the sign $(-1)^{i-1}(-1)(-1)^{i-1} = -1$. Therefore $\det Q_2 < 0$.

For $i = 3$, by switching the 1^{st} and 2^{nd} columns, we have

$$\det Q_3 = \begin{vmatrix} q_{12} & q_{13} & q_{14} & \cdots & q_{1t} \\ q_{22} & q_{23} & q_{24} & \cdots & q_{2t} \\ q_{42} & q_{43} & q_{44} & \cdots & q_{4t} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ q_{t2} & q_{t3} & q_{t4} & \cdots & q_{tt} \end{vmatrix} = (-1) \begin{vmatrix} q_{13} & q_{12} & q_{14} & \cdots & q_{1t} \\ q_{23} & q_{22} & q_{24} & \cdots & q_{2t} \\ q_{43} & q_{42} & q_{44} & \cdots & q_{4t} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ q_{t3} & q_{t2} & q_{t4} & \cdots & q_{tt} \end{vmatrix}.$$

By the case $i = 2$, we have $\det Q_3 > 0$.

.....

For $i = l$,

$$\det Q_l = \begin{vmatrix} q_{12} & \cdots & q_{1(l-1)} & q_{1l} \\ q_{22} & \cdots & q_{2(l-1)} & q_{2l} \\ \vdots & \ddots & \vdots & \vdots \\ q_{(l-1)2} & \cdots & q_{(l-1)(l-1)} & q_{(l-1)l} \end{vmatrix} = (-1)^{l-2} \begin{vmatrix} q_{1l} & q_{12} & \cdots & q_{1(l-1)} \\ q_{2l} & q_{22} & \cdots & q_{2(l-1)} \\ \vdots & \vdots & \ddots & \vdots \\ q_{(l-1)l} & q_{(l-1)2} & \cdots & q_{(l-1)(l-1)} \end{vmatrix}.$$

By the case $i = 2$, the sign of $\det Q_l$ is $(-1)^{l-2}(-1) = (-1)^{l-1}$.

Thus for any $t \geq 2$, the sign of $\det Q_i$ is $(-1)^{i-1}$ for $1 \leq i \leq t$.

Step 2. To show $\det Q_1 = \max_{i=1,2,\dots,t} |\det Q_i|$, i.e.

$$\det Q_1 \geq |\det Q_i|, \quad i = 2, 3, \dots, t.$$

Here we only proof $\det Q_1 \geq |\det Q_2|$ and the rest can be obtained by switching columns.

By Step 1, we know that $\det Q_1 > 0$ and $\det Q_2 < 0$. Thus

$$\begin{aligned} \det Q_1 - |\det Q_2| &= \det Q_1 + \det Q_2 \\ &= \begin{vmatrix} q_{12} + q_{22} & q_{13} + q_{23} & \cdots & q_{1t} + q_{2t} \\ q_{32} & q_{33} & \cdots & q_{3t} \\ \vdots & \vdots & \ddots & \vdots \\ q_{t2} & q_{t3} & \cdots & q_{tt} \end{vmatrix} \\ &= \begin{vmatrix} \bar{q}_2 & \bar{q}_3 & \cdots & \bar{q}_t \\ q_{32} & q_{33} & \cdots & q_{3t} \\ \vdots & \vdots & \ddots & \vdots \\ q_{t2} & q_{t3} & \cdots & q_{tt} \end{vmatrix} = \begin{vmatrix} \bar{q}_2 & q_{23} & \cdots & q_{2t} \\ \bar{q}_3 & q_{33} & \cdots & q_{3t} \\ \vdots & \vdots & \ddots & \vdots \\ \bar{q}_t & q_{t3} & \cdots & q_{tt} \end{vmatrix} \\ &= \bar{q}_2 \det Q'_2 + (-1)\bar{q}_3 \det Q'_3 + \cdots + (-1)^{t-2}\bar{q}_t \det Q'_t, \end{aligned} \quad (3.10)$$

where $Q'_i \in \mathbb{R}^{(t-2) \times (t-2)}$ ($i = 2, 3, \dots, t$) is obtained from eliminating the $(i-1)^{th}$ row of the following $(t-1) \times (t-2)$ matrix

$$Q' := \begin{pmatrix} q_{23} & q_{24} & \cdots & q_{2t} \\ q_{33} & q_{34} & \cdots & q_{3t} \\ q_{43} & q_{44} & \cdots & q_{4t} \\ \vdots & \vdots & \ddots & \vdots \\ q_{t3} & q_{t4} & \cdots & q_{tt} \end{pmatrix}.$$

By Step 1, the sign of $\det Q'_i$ is $(-1)^{i-2}$.

Thus the i^{th} term of (3.10) has the sign $(-1)^{i-1}(-1)^{i-1} = 1$.

Therefore $\det Q_1 - |\det Q_2| > 0$. □

Lemma 3.2 For an integer $t \geq 3$, let $\tilde{Q}_{t \times (t-2)} \in \mathbb{R}^{t \times (t-2)}$ be a matrix as follows

$$\begin{pmatrix} q_{13} & q_{14} & \cdots & q_{1t} \\ q_{23} & q_{24} & \cdots & q_{2t} \\ q_{33} & q_{34} & \cdots & q_{3t} \\ q_{43} & q_{44} & \cdots & q_{4t} \\ \vdots & \vdots & \ddots & \vdots \\ q_{t3} & q_{t4} & \cdots & q_{tt} \end{pmatrix} \quad (3.11)$$

and it satisfies for $i = 1, 2, \dots, t$ and $j = 3, 4, \dots, t$

$$q_{jj} > 0, \quad q_{ij} = q_{ji} < 0 \quad (i \neq j) \quad \text{and} \quad \bar{q}_j := \sum_{i=1}^t q_{ij} > 0. \quad (3.12)$$

Let $\tilde{Q}_{ij} \in \mathbb{R}^{(t-2) \times (t-2)}$ ($1 \leq i < j \leq t$) denote the matrix obtained from eliminating the i^{th} and j^{th} row of the above matrix $\tilde{Q}_{t \times (t-2)}$. Then

$$\det \tilde{Q}_{12} = \max_{1 \leq i < j \leq t} |\det \tilde{Q}_{ij}|.$$

Proof: Consider Q_{1j} to be obtained from eliminating the $(j-1)^{th}$ row of the following $(t-1) \times (t-2)$ matrix

$$\begin{pmatrix} q_{23} & q_{24} & \cdots & q_{2t} \\ q_{33} & q_{34} & \cdots & q_{3t} \\ q_{43} & q_{44} & \cdots & q_{4t} \\ \vdots & \vdots & \ddots & \vdots \\ q_{t3} & q_{t4} & \cdots & q_{tt} \end{pmatrix}.$$

By Lemma 3.1, we have $\det \tilde{Q}_{12} > 0$ and

$$\det \tilde{Q}_{12} \geq |\det \tilde{Q}_{1j}|. \quad (3.13)$$

Similarly, $\det \tilde{Q}_{12} \geq |\det \tilde{Q}_{2j}|$.

In the following we only prove $\det \tilde{Q}_{12} \geq |\det \tilde{Q}_{34}|$ and the rest can be obtained by switching columns.

To estimate $|\det \tilde{Q}_{34}|$ we first compare $\det \tilde{Q}_{34}$ with $\det \tilde{Q}_{14}$ and $\det \tilde{Q}_{13}$.

$$\begin{aligned}
\det \tilde{Q}_{34} - \det \tilde{Q}_{14} &= \begin{vmatrix} q_{13} + q_{33} & q_{14} + q_{34} & q_{15} + q_{35} & \cdots & q_{1t} + q_{3t} \\ q_{23} & q_{24} & q_{25} & \cdots & q_{2t} \\ q_{53} & q_{54} & q_{55} & \cdots & q_{5t} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ q_{t3} & q_{t4} & q_{t5} & \cdots & q_{tt} \end{vmatrix} \\
&= \begin{vmatrix} q_{31} + q_{33} & q_{32} & q_{35} & \cdots & q_{3t} \\ q_{41} + q_{43} & q_{42} & q_{45} & \cdots & q_{4t} \\ q_{51} + q_{53} & q_{52} & q_{55} & \cdots & q_{5t} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ q_{t1} + q_{t3} & q_{t2} & q_{t5} & \cdots & q_{tt} \end{vmatrix} = \begin{vmatrix} q_{31} + q_{33} + \sum_{j=5}^t \lambda_j q_{3j} & q_{32} & q_{35} & \cdots & q_{3t} \\ q_{41} + q_{43} + \sum_{j=5}^t \lambda_j q_{4j} & q_{42} & q_{45} & \cdots & q_{4t} \\ q_{51} + q_{53} + \sum_{j=5}^t \lambda_j q_{5j} & q_{52} & q_{55} & \cdots & q_{5t} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ q_{t1} + q_{t3} + \sum_{j=5}^t \lambda_j q_{tj} & q_{t2} & q_{t5} & \cdots & q_{tt} \end{vmatrix} \\
&= (q_{31} + q_{33} + \sum_{j=5}^t \lambda_j q_{3j}) \det \tilde{Q}'_3 + (-1)(q_{41} + q_{43} + \sum_{j=5}^t \lambda_j q_{4j}) \det \tilde{Q}'_4 \\
&\quad + (-1)^2(q_{51} + q_{53} + \sum_{j=5}^t \lambda_j q_{5j}) \det \tilde{Q}'_5 + \cdots + (-1)^{t-3}(q_{t1} + q_{t3} + \sum_{j=5}^t \lambda_j q_{tj}) \det \tilde{Q}'_t, \quad (3.14)
\end{aligned}$$

where $\tilde{Q}'_i (i = 3, 4, \dots, t)$ is the matrix obtained from eliminating the $(i-2)^{th}$ row of the following matrix

$$\begin{pmatrix} q_{32} & q_{35} & \cdots & q_{3t} \\ q_{42} & q_{45} & \cdots & q_{4t} \\ q_{52} & q_{55} & \cdots & q_{5t} \\ \vdots & \vdots & \ddots & \vdots \\ q_{t2} & q_{t5} & \cdots & q_{tt} \end{pmatrix}, \quad (3.15)$$

and for $j = 5, 6, \dots, t$,

$$\lambda_j = \begin{cases} 0 & \text{if } (-1)^{j-3} \det \tilde{Q}'_j \geq 0, \\ 1 & \text{if } (-1)^{j-3} \det \tilde{Q}'_j < 0. \end{cases} \quad (3.16)$$

By Lemma 3.1 $\det \tilde{Q}'_3 < 0$, $\det \tilde{Q}'_4 < 0$ and

$$q_{31} + q_{33} + \sum_{j=5}^t \lambda_j q_{3j} > \bar{q}_3 > 0, \quad q_{41} + q_{43} + \sum_{j=5}^t \lambda_j q_{4j} < 0.$$

Thus, the first two terms of (3.14) are both negative.

By the definition of λ_i in (3.16),

i) if $(-1)^{i-3} \det \tilde{Q}'_i \geq 0$, $\lambda_i = 0$, then

$$q_{i1} + q_{i3} + \sum_{j=5}^t \lambda_j q_{ij} = q_{i1} + q_{i3} + \sum_{j \neq i, j=5}^t \lambda_j q_{ij} < 0;$$

ii) if $(-1)^{i-3} \det \tilde{Q}'_i < 0$, $\lambda_i = 1$, then

$$q_{i1} + q_{i3} + \sum_{j=5}^t \lambda_j q_{ij} > \bar{q}_i > 0.$$

Thus, the $(i-2)^{th}$ ($i = 5, 6, \dots, t$) term of (3.14)

$$(-1)^{i-3} (q_{i1} + q_{i3} + \sum_{j=5}^t \lambda_j q_{ij}) \det \tilde{Q}'_i \leq 0.$$

Therefore, $\det \tilde{Q}_{34} - \det \tilde{Q}_{14} < 0$, i.e. $\det \tilde{Q}_{34} < \det \tilde{Q}_{14}$.

Similarly,

$$\begin{aligned} \det \tilde{Q}_{34} - \det \tilde{Q}_{13} &= \begin{vmatrix} q_{13} + q_{43} & q_{14} + q_{44} & q_{15} + q_{45} & \cdots & q_{1t} + q_{4t} \\ q_{23} & q_{24} & q_{25} & \cdots & q_{2t} \\ q_{53} & q_{54} & q_{55} & \cdots & q_{5t} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ q_{t3} & q_{t4} & q_{t5} & \cdots & q_{tt} \end{vmatrix} \\ &= \begin{vmatrix} q_{31} + q_{34} & q_{32} & q_{35} & \cdots & q_{3t} \\ q_{41} + q_{44} & q_{42} & q_{45} & \cdots & q_{4t} \\ q_{51} + q_{54} & q_{52} & q_{55} & \cdots & q_{5t} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ q_{t1} + q_{t4} & q_{t2} & q_{t5} & \cdots & q_{tt} \end{vmatrix} = \begin{vmatrix} q_{31} + q_{34} + \sum_{j=5}^t \lambda'_j q_{3j} & q_{32} & q_{35} & \cdots & q_{3t} \\ q_{41} + q_{44} + \sum_{j=5}^t \lambda'_j q_{4j} & q_{42} & q_{45} & \cdots & q_{4t} \\ q_{51} + q_{54} + \sum_{j=5}^t \lambda'_j q_{5j} & q_{52} & q_{55} & \cdots & q_{5t} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ q_{t1} + q_{t4} + \sum_{j=5}^t \lambda'_j q_{tj} & q_{t2} & q_{t5} & \cdots & q_{tt} \end{vmatrix} \\ &= (q_{31} + q_{34} + \sum_{j=5}^t \lambda'_j q_{3j}) \det \tilde{Q}'_3 + (-1)(q_{41} + q_{44} + \sum_{j=5}^t \lambda'_j q_{4j}) \det \tilde{Q}'_4 \\ &\quad + (-1)^2 (q_{51} + q_{54} + \sum_{j=5}^t \lambda'_j q_{5j}) \det \tilde{Q}'_5 + \cdots + (-1)^{t-3} (q_{t1} + q_{t4} + \sum_{j=5}^t \lambda'_j q_{tj}) \det \tilde{Q}'_t, \quad (3.17) \end{aligned}$$

where \tilde{Q}'_i ($i = 3, 4, \dots, t$) is the same matrix obtained from eliminating the $(i-2)^{th}$ row of the matrix (3.15) and for $j = 5, 6, \dots, t$,

$$\lambda'_j = \begin{cases} 1 & \text{if } (-1)^{j-3} \det \tilde{Q}'_j \geq 0, \\ 0 & \text{if } (-1)^{j-3} \det \tilde{Q}'_j < 0. \end{cases} \quad (3.18)$$

By Lemma 3.1 $\det \tilde{Q}'_3 < 0$, $\det \tilde{Q}'_4 < 0$ and

$$q_{31} + q_{34} + \sum_{j=5}^t \lambda'_j q_{3j} < 0, \quad q_{41} + q_{44} + \sum_{j=5}^t \lambda'_j q_{4j} > \bar{q}_4 > 0.$$

Thus, the first two terms of (3.17) are both positive.

By the definition of λ'_i in (3.18),

i) if $(-1)^{i-3} \det \tilde{Q}'_i \geq 0$, $\lambda'_i = 1$, then

$$q_{i1} + q_{i4} + \sum_{j=5}^t \lambda'_j q_{ij} > \bar{q}_i > 0;$$

ii) if $(-1)^{i-3} \det \tilde{Q}'_i < 0$, $\lambda'_i = 0$, then

$$q_{i1} + q_{i3} + \sum_{j=5}^t \lambda_j q_{ij} = q_{i1} + q_{i4} + \sum_{j \neq i, j=5}^t \lambda'_j q_{ij} < 0.$$

Thus, the $(i-2)^{th}$ ($i = 5, 6, \dots, t$) term of (3.17)

$$(-1)^{i-3} (q_{i1} + q_{i4} + \sum_{j=5}^t \lambda'_j q_{ij}) \det \tilde{Q}'_i \geq 0.$$

Therefore, $\det \tilde{Q}_{34} - \det \tilde{Q}_{13} > 0$, i.e. $\det \tilde{Q}_{34} > \det \tilde{Q}_{13}$.

Hence, by (3.13),

$$|\det \tilde{Q}_{34}| < \max(|\det \tilde{Q}_{14}|, |\det \tilde{Q}_{13}|) < \det \tilde{Q}_{12}.$$

We complete the proof. □

Lemma 3.3 For an integer $t \geq 1$, let $Q_{t \times t} \in \mathbb{R}^{t \times t}$ be a matrix as follows

$$\begin{pmatrix} q_{11} & q_{12} & \cdots & q_{1t} \\ q_{21} & q_{22} & \cdots & q_{2t} \\ \vdots & \vdots & \ddots & \vdots \\ q_{t1} & q_{t2} & \cdots & q_{tt} \end{pmatrix} \quad (3.19)$$

and it satisfies for $i, j = 1, 2, \dots, t$

$$q_{ii} > 0, \quad q_{ij} = q_{ji} < 0 \quad (i \neq j) \quad \text{and} \quad \bar{q}_i := \sum_{j=1}^t q_{ij} > 0. \quad (3.20)$$

Then we have

$$\bar{q}_1 \begin{vmatrix} q_{22} & \cdots & q_{2t} \\ \vdots & \ddots & \vdots \\ q_{t2} & \cdots & q_{tt} \end{vmatrix} \leq \det Q \leq \sum_{i,j=1}^t q_{ij} \begin{vmatrix} q_{22} & \cdots & q_{2t} \\ \vdots & \ddots & \vdots \\ q_{t2} & \cdots & q_{tt} \end{vmatrix}.$$

Proof: We have

$$\begin{aligned} \det Q &= \begin{vmatrix} \bar{q}_1 & q_{12} & \cdots & q_{1t} \\ \bar{q}_2 & q_{22} & \cdots & q_{2t} \\ \vdots & \vdots & \ddots & \vdots \\ \bar{q}_t & q_{t2} & \cdots & q_{tt} \end{vmatrix} \\ &= \bar{q}_1 \det Q_1 + (-1)\bar{q}_2 \det Q_2 + \cdots + (-1)^{t-1} \bar{q}_t \det Q_t, \end{aligned}$$

where $Q_i (i = 1, 2, \dots, t)$ is the matrix obtained from eliminating the i^{th} row of the following matrix

$$\begin{pmatrix} q_{12} & q_{13} & \cdots & q_{1t} \\ q_{22} & q_{23} & \cdots & q_{2t} \\ q_{32} & q_{33} & \cdots & q_{3t} \\ \vdots & \vdots & \ddots & \vdots \\ q_{t2} & q_{t3} & \cdots & q_{tt} \end{pmatrix}.$$

By Lemma 3.1, the i^{th} term has the sign $(-1)^{i-1}(-1)^{i-1} = 1$. Thus each term are positive.

Therefore

$$\det Q \geq 1^{st} \text{ term} = \bar{q}_1 \begin{vmatrix} q_{22} & \cdots & q_{2t} \\ \vdots & \ddots & \vdots \\ q_{t2} & \cdots & q_{tt} \end{vmatrix}$$

and

$$\det Q \leq \sum_{i=1}^t \bar{q}_i \begin{vmatrix} q_{22} & \cdots & q_{2t} \\ \vdots & \ddots & \vdots \\ q_{t2} & \cdots & q_{tt} \end{vmatrix} = \sum_{i,j=1}^t q_{ij} \begin{vmatrix} q_{22} & \cdots & q_{2t} \\ \vdots & \ddots & \vdots \\ q_{t2} & \cdots & q_{tt} \end{vmatrix}.$$

□

Proof of Theorem 3.1: We only estimate $|\alpha_1 - \alpha_2|$ and the rest can be obtained by switching columns.

By the system (3.5), we can get the formula for the difference of α_1 and α_2 :

$$\alpha_1 - \alpha_2 = \frac{\begin{vmatrix} \beta_1 & p_{11} + p_{12} & p_{13} & \cdots & p_{1s} \\ \beta_2 & p_{21} + p_{22} & p_{23} & \cdots & p_{2s} \\ \beta_3 & p_{31} + p_{32} & p_{33} & \cdots & p_{3s} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \beta_s & p_{s1} + p_{s2} & p_{s3} & \cdots & p_{ss} \end{vmatrix}}{\begin{vmatrix} p_{11} & p_{12} & \cdots & p_{1s} \\ p_{21} & p_{22} & \cdots & p_{2s} \\ \vdots & \vdots & \ddots & \vdots \\ p_{s1} & p_{s2} & \cdots & p_{ss} \end{vmatrix}} = \frac{\begin{vmatrix} \beta_1 & \bar{p}_1 & p_{13} & \cdots & p_{1s} \\ \beta_2 & \bar{p}_2 & p_{23} & \cdots & p_{2s} \\ \beta_3 & \bar{p}_3 & p_{33} & \cdots & p_{3s} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \beta_s & \bar{p}_s & p_{s3} & \cdots & p_{ss} \end{vmatrix}}{\begin{vmatrix} p_{11} & p_{12} & \cdots & p_{1s} \\ p_{21} & p_{22} & \cdots & p_{2s} \\ \vdots & \vdots & \ddots & \vdots \\ p_{s1} & p_{s2} & \cdots & p_{ss} \end{vmatrix}} := \frac{\det \tilde{P}}{\det P}.$$

We estimate the determinants of the above two matrices separately.

By Lemma 3.2 and condition (3.4), we have

$$|\det \tilde{P}| \leq C \begin{vmatrix} p_{33} & \cdots & p_{3s} \\ \vdots & \ddots & \vdots \\ p_{s3} & \cdots & p_{ss} \end{vmatrix},$$

where $C := C(C_0)$ is a positive constant.

By Lemma 3.3 and condition (3.4), we have

$$\begin{aligned} \det P &\geq \bar{p}_1 \begin{vmatrix} p_{22} & \cdots & p_{2s} \\ \vdots & \ddots & \vdots \\ p_{s2} & \cdots & p_{ss} \end{vmatrix} \geq \bar{p}_1 \sum_{j=2}^s p_{2j} \begin{vmatrix} p_{33} & \cdots & p_{3s} \\ \vdots & \ddots & \vdots \\ p_{s3} & \cdots & p_{ss} \end{vmatrix} \\ &= \bar{p}_1 (\bar{p}_2 - p_{12}) \begin{vmatrix} p_{33} & \cdots & p_{3s} \\ \vdots & \ddots & \vdots \\ p_{s3} & \cdots & p_{ss} \end{vmatrix} \geq \frac{1}{C_0} |p_{12}| \begin{vmatrix} p_{33} & \cdots & p_{3s} \\ \vdots & \ddots & \vdots \\ p_{s3} & \cdots & p_{ss} \end{vmatrix}. \end{aligned}$$

Therefore,

$$|\alpha_1 - \alpha_2| = \frac{|\det \tilde{P}|}{|\det P|} \leq \frac{C}{|p_{12}|},$$

where $C := C(C_0)$ is a positive constant. \square

3.1.2 Proof of Theorem 1.5

Similar to the two inclusions case in chapter 2, we decompose u into $m + 1$ parts:

$$u = v_0 + \sum_{i=1}^m C_i \cdot v_i, \quad (3.21)$$

where $v_i \in H^1(\tilde{\Omega})$ ($i = 0, 1, 2, \dots, m$), are determined by the following equations:

for $i = 0$,

$$\begin{cases} \Delta v_0 = 0 & \text{in } \tilde{\Omega}, \\ v_0 = 0 & \text{on } \partial D_1, \partial D_2, \dots, \partial D_m, \\ v_0 = \varphi & \text{on } \partial \Omega. \end{cases} \quad (3.22)$$

for $i = 1, 2, \dots, m$,

$$\begin{cases} \Delta v_i = 0 & \text{in } \tilde{\Omega}, \\ v_i = 1 & \text{on } \partial D_i, \\ v_i = 0 & \text{on } \partial D_j, \text{ for } j \neq i, \\ v_i = 0 & \text{on } \partial \Omega; \end{cases} \quad (3.23)$$

Since u satisfies the integral conditions in equation (3.1), using the decomposition formula (3.21), we see that the vector (C_1, C_2, \dots, C_m) is uniquely determined by the system of linear equations

$$\begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1m} \\ a_{21} & a_{22} & \cdots & a_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mm} \end{pmatrix} \cdot \begin{pmatrix} C_1 \\ C_2 \\ \vdots \\ C_m \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{pmatrix} \quad (3.24)$$

where

$$a_{ij} := \int_{\partial D_j} \frac{\partial v_i}{\partial \nu}, \quad (i, j = 1, 2, \dots, m), \quad (3.25)$$

$$b_i := \int_{\partial D_i} \frac{\partial v_0}{\partial \nu}, \quad (i = 1, 2, \dots, m). \quad (3.26)$$

Similar to the two inclusions case in chapter 2, we investigate some properties of v_i ($i = 0, 1, \dots, m$), the matrix A and the vector b in equation (3.24). Here we state several lemmas without proofs since the proofs are same as those of Lemma 2.1, 2.3 and 2.4 in chapter 2.

Lemma 3.4 *Let v_0, v_1, \dots, v_m be defined by equations (3.22) and (3.23), then for $n \geq 2$ and for $i = 1, 2, \dots, m$ we have*

$$\|\nabla v_0\|_{L^\infty(\tilde{\Omega})} \leq C, \quad \|\nabla v_i\|_{L^\infty(\tilde{\Omega})} \leq \frac{C}{\varepsilon}, \quad \left\| \frac{\partial v_i}{\partial \nu} \right\|_{L^\infty(\partial\Omega)} \leq C.$$

Lemma 3.5 *For $1 \leq i, j \leq m$, let a_{ij} and b_i be defined as in (3.25) and (3.26), then they satisfy the following:*

$$1) \quad a_{ii} < 0, \quad a_{ij} = a_{ji} > 0 \quad (i \neq j),$$

$$2) \quad -C \leq \sum_{1 \leq j \leq m} a_{ij} \leq -\frac{1}{C},$$

$$3) \quad |b_i| \leq C,$$

where $C > 0$ is a universal constant.

Remark 3.1 *From property (1) and (2) in Lemma 3.5, we can prove that the matrix A is nonsingular, it serves a good exercise in linear algebra.*

Next, we derive estimates of the order of a_{ij} with respect to ε_{ij} .


Lemma 3.6 *Let a_{ij} be defined as in (3.25), there exists a universal constant $C > 0$, such that for any $1 \leq i \neq j \leq m$,*

$$\begin{aligned} -\frac{C}{\sqrt{\min_{j \neq i} \varepsilon_{ij}}} &< a_{ii} < -\frac{1}{C \sqrt{\min_{j \neq i} \varepsilon_{ij}}}, & \frac{1}{\sqrt{\varepsilon_{ij}}} &< a_{ij} < \frac{C}{\sqrt{\varepsilon_{ij}}}, & \text{for } n = 2, \\ -C |\ln(\min_{j \neq i} \varepsilon_{ij})| &< a_{ii} < -\frac{1}{C} |\ln(\min_{j \neq i} \varepsilon_{ij})|, & \frac{1}{C} \ln \varepsilon_{ij} &< a_{ij} < C \ln \varepsilon_{ij}, & \text{for } n = 3, \\ -C &< a_{ii} < -\frac{1}{C}, & \frac{1}{C} &< a_{ij} < C, & \text{for } n \geq 4. \end{aligned}$$

Proof: Without loss of generality, we assume $i = 1, j = 2$. The proof of the estimate about a_{11} is the same as that in Lemma 2.5, 2.6, and 2.7. Here we prove the estimate about a_{12} . Let Γ be a $C^{2,\alpha}$ closed simple curve such that D_2 is inside the area bounded by the curve. Denote:

$$\Gamma_1 := \Gamma \cap D_1 \quad \Gamma_2 := \Gamma \cap (\cup_{2 \leq i \leq m} \partial D_i) \quad \Gamma_3 := \Gamma \setminus (\Gamma_1 \cup \Gamma_2).$$

It's easy to draw Γ such that Γ_1 contains the closest point of D_1 to D_2 , and $\|\nabla v_1\|_{L^\infty(\Gamma_3)} < C$ for some universal positive constant C . We can also show that $\|\nabla v_1\|_{L^\infty(\Gamma_2)} < C$ by comparison principle. Look at the following figure for example.



By the harmonicity of v_1 , we have

$$\begin{aligned}
 a_{12} &= \int_{\partial D_2} \frac{\partial v_1}{\partial \nu} = \int_{\Gamma} \frac{\partial v_1}{\partial \nu} \\
 &= \int_{\Gamma_1} \frac{\partial v_1}{\partial \nu} + \int_{\Gamma_2} \frac{\partial v_1}{\partial \nu} + \int_{\Gamma_3} \frac{\partial v_1}{\partial \nu} \\
 &= \int_{\Gamma_1} \frac{\partial v_1}{\partial \nu} + O(1).
 \end{aligned} \tag{3.27}$$

The last equality of (3.27) is from the fact that $\int_{\Gamma_2} \frac{\partial v_1}{\partial \nu} \leq C$ and $\|\nabla v_1\|_{L^\infty(\Gamma_3)} < C$.

Using the above facts again, it follows that

$$\int_{\Gamma_1} \frac{\partial v_1}{\partial \nu} = \int_{\Gamma} v_1 \frac{\partial v_1}{\partial \nu} + O(1) = \int_{\widetilde{D_2}} |\nabla v_1|^2 + O(1), \tag{3.28}$$

where $\widetilde{D_2}$ is denoted as the region bounded by Γ and ∂D_2 .

Therefore, by combining (3.27) and (3.28), we have

$$a_{12} = \int_{\widetilde{D_2}} |\nabla v_1|^2 + O(1).$$

Using the same method as that in the proofs of Lemma 2.5, 2.6, and 2.7, we have

$$\begin{aligned} \frac{1}{C\sqrt{\varepsilon_{12}}} &< \int_{\widetilde{D_2}} |\nabla v_1|^2 < \frac{C}{\sqrt{\varepsilon_{12}}}, & \text{for } n = 2 \\ \frac{1}{C} |\ln \varepsilon_{12}| &< \int_{\widetilde{D_2}} |\nabla v_1|^2 < C |\ln \varepsilon_{12}|, & \text{for } n = 3 \\ \frac{1}{C} &< \int_{\widetilde{D_2}} |\nabla v_1|^2 < C, & \text{for } n \geq 4. \end{aligned}$$

Therefore,

$$\begin{aligned} \frac{1}{\sqrt{\varepsilon_{12}}} &< a_{12} < \frac{C}{\sqrt{\varepsilon_{12}}}, & \text{for } n = 2, \\ \frac{1}{C} \ln \varepsilon_{12} &< a_{12} < C \ln \varepsilon_{12}, & \text{for } n = 3, \\ \frac{1}{C} &< a_{12} < C, & \text{for } n \geq 4. \end{aligned}$$

□

Now Applying Theorem 3.1 and Lemma 3.5, 3.6, we have

Proposition 3.1 *Let $u \in H^1(\Omega) \cap C^1(\overline{\Omega})$ be the solution to equation (3.1) and let C_i be the value of u on D_i , there exists a universal constant $C > 0$ which depends only on $n, \kappa_0, r_0, \|\partial\Omega\|_{C^{2,\alpha}}, \{\|\partial D_i\|_{C^{2,\alpha}}\}$, but independent of $\{\varepsilon_{ij}\}$ such that*

$$\begin{aligned} |C_i - C_j| &\leq C\sqrt{\varepsilon_{ij}} & \text{for } n = 2, \\ |C_i - C_j| &\leq C \frac{1}{|\ln \varepsilon_{ij}|} & \text{for } n = 3, \\ |C_i - C_j| &\leq C & \text{for } n \geq 4. \end{aligned} \tag{3.29}$$

Now we are ready to complete the proof of Theorem 1.5.

Proof of Theorem 1.5 We prove the estimates in dimension 2, the higher dimension cases follow from the same idea. Without loss of generality, we assume $x_{ij}^0 = 0$ and $C_i \leq C_j$. Since $\Omega, D_1, D_2, \dots, D_m$ satisfy the properties listed in (1.19), we can choose δ small enough such that if the distance ε_{ij} of D_i and D_j satisfies $\varepsilon_{ij} < \delta$, then the ball $B(0, 4\delta)$ only intersects with D_i and D_j .

By the maximum principle,

$$\|u\|_{L^\infty(\tilde{\Omega})} \leq \|\varphi\|_{L^\infty(\partial\Omega)}.$$

Using gradient estimates and boundary estimates of harmonic function, we immediately see that, for some universal constant $C > 0$,

$$\|\nabla u\|_{L^\infty(\tilde{\Omega} \cap B(0, 3\delta) \setminus \overline{B(0, \delta)})} \leq C\|u\|_{L^\infty(\tilde{\Omega})} \leq C\|\varphi\|_{L^\infty(\partial\Omega)}. \quad (3.30)$$

Since $\varepsilon_{ij} < \delta$, we have

$$B(0, \delta) \subset B(x^i, 2\delta) \subset B(0, 3\delta). \quad (3.31)$$

Therefore we can find a harmonic function $w \in C^2(B(x^i, 2\delta) \setminus \overline{D_i}) \cap C^0(\overline{B(x^i, 2\delta) \setminus D_i})$ satisfying the following boundary conditions

- $w = 0$ on $B(x^i, \delta) \cap \partial D_i$,
- $w > |u - C_i|$ on $\partial B(x^i, 2\delta) \setminus (D_i \cup D_j)$, (This is from (3.30) and (3.31))
- $w > 0$ on $B(x^i, 2\delta) \setminus \overline{D_i}$.

By the Hopf Lemma, we know

$$\frac{\partial w}{\partial \nu} > 0 \quad \text{on } B(x^i, \delta) \cap \partial D_i.$$

Then we can choose α_0 small enough such that if $\varepsilon_{ij} < \alpha_0$, then

$$w > C\varepsilon_{ij} \quad \text{on } B(x^j, \alpha_0) \cap \partial D_j, \quad (3.32)$$

for some universal constant $C > 0$.

Now we Claim: there exists a universal constant $C > 0$ such that

$$\frac{Cw}{\sqrt{\varepsilon_{ij}}} \geq u - C_i \geq -\frac{Cw}{\sqrt{\varepsilon_{ij}}} \quad \text{on } B(x^i, 2\delta) \setminus \overline{D_i \cup D_j}. \quad (3.33)$$

Proof of the Claim: By the maximum principle, we only need to show that there exists some universal constant $C > 0$ such that

$$\frac{Cw}{\sqrt{\varepsilon_{ij}}} \geq u - C_i \geq -\frac{Cw}{\sqrt{\varepsilon_{ij}}} \quad \text{on } \partial(B(x^i, 2\delta) \setminus \overline{D_i \cup D_j}).$$

According to our construction of w , we know that $w = 0$ on $B(x^i, \delta) \cap \partial D_i$, and $w > |u - C_i|$ on $\partial B(x^i, 2\delta) \setminus (D_i \cup D_j)$, therefore, we only need to show that we can find some universal constant $C > 0$ such that

$$\frac{Cw}{\sqrt{\varepsilon_{ij}}} \geq |u - C_i| \quad \text{on } \partial D_j \cap B(x^i, 2\delta).$$

According to (3.32) and Proposition 3.1, we have

$$\frac{Cw}{\sqrt{\varepsilon_{ij}}} \geq C\sqrt{\varepsilon_{ij}} \geq |C_j - C_i| = |u - C_i| \quad \text{on } B(x^j, \alpha_0) \cap \partial D_j.$$

Since $w \geq C$ for some universal constant $C > 0$ on $\partial(D_j \cap B(x^i, 2\delta)) \setminus B(x^j, \alpha_0)$, by choosing C large enough, we can make

$$\frac{Cw}{\sqrt{\varepsilon_{ij}}} \geq |u - C_i| \quad \text{on } \partial D_j \cap B(x^i, 2\delta).$$

Therefore our claim is proved.

By the comparison principle, according to (3.33), we see that

$$\frac{C}{\sqrt{\varepsilon_{ij}}} \geq \frac{\partial}{\partial \nu} \frac{Cw}{\sqrt{\varepsilon_{ij}}} \geq \left| \frac{\partial u}{\partial \nu} \right| = |\nabla u| \quad \text{on } B(x^i, \delta) \cap \partial D_i.$$

Applying the same method, we can get for some universal constant $\delta > 0$,

$$\frac{C}{\sqrt{\varepsilon_{ij}}} \geq \left| \frac{\partial u}{\partial \nu} \right| = |\nabla u| \quad \text{on } B(x^j, \delta) \cap \partial D_j.$$

Therefore by the maximum principle, we have

$$\|\nabla u\|_{L^\infty(\tilde{\Omega} \cap B(0, \delta))} \leq \frac{C}{\sqrt{\varepsilon_{ij}}} \|\varphi\|_{C^2(\partial\Omega)}.$$

□

For the equations with more general coefficients, with the same proofs in section 2.3 we can also derive Theorem 1.5.

3.2 The insulated conductivity problem

As stated in the introduction, for the insulated conductivity problem, we only need to consider the local situation (1.24). Since the proof for the case with multiple inclusions is

same as the one with two inclusions, here we assume that there are only two conductors D_1 and D_2 . We consider harmonic equation first and the equation with general coefficients can be obtained similarly. In this section we use, unless otherwise stated, C to denote various positive constants whose values may change from line to line and which depend only on n , κ_0 , r_0 , $\|\partial\Omega\|_{C^{2,\alpha}}$, $\|\partial D_1\|_{C^{2,\alpha}}$ and $\|\partial D_2\|_{C^{2,\alpha}}$.

The insulated conductivity problem can be described as the following local version

$$\begin{cases} \Delta u = 0 & \text{in } \tilde{\Omega} \cap B(x_{12}^0, 2\delta), \\ \frac{\partial u}{\partial \nu}|_+ = 0 & \text{on } (\partial D_1 \cup \partial D_2) \cap B(x_{12}^0, 2\delta), \\ |u| \leq 1 & \text{in } \tilde{\Omega} \cap B(x_{12}^0, 2\delta), \end{cases} \quad (3.34)$$

where $B(x_{12}^0, 2\delta)$ is defined in (1.20).

Without loss of generality, by a translation and rotation of the axis, we may also assume that D_1, D_2 are two strictly convex subdomains in $\Omega \subset \mathbb{R}^n$ which satisfy the following:

$$(-\varepsilon/2, 0') \in \partial D_1, \quad (\varepsilon/2, 0') \in \partial D_2, \quad \varepsilon = \text{dist}(\partial D_1, \partial D_2) = \text{dist}(D_1, D_2). \quad (3.35)$$

Near the origin, we can find a ball $B(0, r)$ such that the portion of ∂D_i ($i = 1, 2$) in $B(0, r)$ is strictly convex, where $r > 0$ is independent of ε . Then $\partial D_1 \cap B(0, r)$ and $\partial D_2 \cap B(0, r)$ can be represented by the graph of $x_1 = -\frac{\varepsilon}{2} - f(x')$ and $x_1 = \frac{\varepsilon}{2} + g(x')$ respectively, where $x' = (x_2, \dots, x_n)$. Thus $f(0') = g(0') = 0$, $\nabla f(0') = \nabla g(0') = 0$, and $\frac{1}{C}I \leq (D^2 f(0')), (D^2 g(0')) \leq CI$.

Proposition 3.2 *Under the hypotheses of Theorem 1.6, let u be the solution of equation (3.34). There exists a positive constants C , such that, for sufficiently small $\varepsilon > 0$,*

$$|\nabla u(x)| \leq \frac{C}{\sqrt{\varepsilon}}, \quad (3.36)$$

if $-\frac{\varepsilon}{2} - f(x') \leq x_1 \leq \frac{\varepsilon}{2} + g(x')$, and $|x'| \leq \frac{\sqrt{\varepsilon}}{2}$.

Before proving the above proposition, we first show the following lemmas.

Lemma 3.7 For any small $\delta > 0$, let $\mathcal{O}_z \doteq \{z \in \mathbb{R}^n \mid |z_1| \leq \delta \text{ and } |z'| \leq 1\}$, and let $w \in H^1(\mathcal{O}_z)$ satisfied the following conditions

$$\partial_{z_j} \left(a^{ij}(z) \partial_{z_i} w \right) = 0 \quad \text{in } \mathcal{O}_z, \quad (3.37)$$

$$a^{1\alpha}(z) = a^{\alpha 1}(z) = 0 \quad \text{on } |z_1| = \delta, \alpha = 2, 3, \dots, n, \quad (3.38)$$

$$\frac{\partial w}{\partial z_1} = 0 \quad \text{on } |z_1| = \delta, \quad (3.39)$$

$$|w| \leq 1 \quad \text{in } \mathcal{O}_z, \quad (3.40)$$

where the $n \times n$ matrix $A(z) := (a^{ij}(z))$ in \mathcal{O}_z is symmetric, $a^{ij}(z) \in C^1(\overline{\mathcal{O}_z})$ and \exists constants $\Lambda \geq \lambda > 0$ such that, for all $\xi \in \mathbb{R}^n$,

$$\lambda |\xi|^2 \leq a^{ij}(z) \xi_i \xi_j \leq \Lambda |\xi|^2 \quad (\forall z \in \mathcal{O}_z).$$

Then there exists a constant $C > 0$ which depends on $n, \lambda, \Lambda, \|a^{ij}(z)\|_{C^1(\overline{\mathcal{O}_z})}$, and is independent of δ , such that

$$\|\nabla w\|_{L^\infty(\frac{1}{2}\mathcal{O}_z)} \leq C,$$

where $\frac{1}{2}\mathcal{O}_z \doteq \{z \in \mathbb{R}^n \mid |z_1| \leq \delta \text{ and } |z'| \leq \frac{1}{2}\}$.

Proof: We construct a new function \tilde{w} by “flipping” w about the hyperplane $z_1 = \delta$ as follows

$$\tilde{w}(z) = \begin{cases} w(z) & \text{if } z \in \mathcal{O}_z, \\ w(2\delta - z_1, z') & \text{if } \delta \leq z_1 \leq 3\delta \text{ and } |z'| \leq 1. \end{cases} \quad (3.41)$$

Thus \tilde{w} extends w to $\{z \in \mathbb{R}^n \mid -\delta \leq z_1 \leq 3\delta \text{ and } |z'| \leq 1\}$. Keep flipping \tilde{w} about $z_1 = -\delta$, $z_1 = 3\delta$, $z_1 = -5\delta$, ..., we get a new function, still denoted as \tilde{w} , which is defined in $\{z \in \mathbb{R}^n \mid |z_1| \leq 1 \text{ and } |z'| \leq 1\}$.

Since w satisfies the condition (3.39) and (3.40), \tilde{w} satisfies

$$\begin{aligned} \partial_{z_j} (\tilde{a}^{ij}(z) \partial_{z_i} \tilde{w}) &= 0 \quad \text{in } \{z \in \mathbb{R}^n \mid |z_1| \leq 1 \text{ and } |z'| \leq 1\}, \\ |\tilde{w}| &\leq 1 \quad \text{in } \{z \in \mathbb{R}^n \mid |z_1| \leq 1 \text{ and } |z'| \leq 1\}, \end{aligned} \quad (3.42)$$

where for $\alpha = 2, 3, \dots, n$,

$$\tilde{a}^{\alpha 1}(z) = \tilde{a}^{1\alpha}(z) = \begin{cases} a^{1\alpha}(z) & \text{if } z \in \mathcal{O}_z, \\ -a^{1\alpha}(2\delta - z_1, z') & \text{if } \delta \leq z_1 \leq 3\delta \text{ and } |z'| \leq 1. \end{cases} \quad (3.43)$$

for other indices

$$\tilde{a}^{ij}(z) = \begin{cases} a^{ij}(z) & \text{if } z \in \mathcal{O}_z, \\ a^{ij}(2\delta - z_1, z') & \text{if } \delta \leq z_1 \leq 3\delta \text{ and } |z'| \leq 1. \end{cases} \quad (3.44)$$

Since $a^{ij}(z) \in C^1(\overline{\mathcal{O}_z})$ satisfy (3.38), $\tilde{a}^{ij}(z)$ is Lipschitz continuous in $\{z \in \mathbb{R}^n \mid |z_1| \leq 1 \text{ and } |z'| \leq 1\}$. Thus, by (3.42) there exists a constant $C > 0$ which depends on n, λ, Λ , $\|a^{ij}(z)\|_{C^1(\overline{\mathcal{O}_z})}$, and is independent of δ , such that

$$|\nabla \tilde{w}(z)| \leq C \quad \text{if } |z_1| \leq \frac{1}{2} \text{ and } |z'| \leq \frac{1}{2}.$$

Since $\tilde{w}(z) = w(z)$ in \mathcal{O}_z , therefore $|\nabla w| \leq C$ in $\frac{1}{2}\mathcal{O}_z$. \square

Lemma 3.8 *For any $\delta, r > 0$, let $\mathcal{O}_y(\delta, r) \doteq \{y \in \mathbb{R}^n \mid |y_1| \leq \delta(1 + |y'|^2) \text{ and } |y'| \leq r\}$, $\mathcal{O}_z(\delta, r) \doteq \{z \in \mathbb{R}^n \mid |z_1| \leq \delta \text{ and } |z'| \leq r\}$. There exist a constant $\bar{r} > 0$ which is independent of δ and a transformation $y = \Phi(z) : \mathcal{O}_z(\delta, r) \longrightarrow \mathcal{O}_y(\delta, r)$, such that $\forall 0 < r \leq \bar{r}$ and $\forall \delta > 0$ sufficient small,*

$$\Phi(\pm\delta, z') = (\pm\delta(1 + |z'|^2), z'), \quad (3.45)$$

i.e. it maps the boundary $\Gamma_z := \{z \in \mathbb{R}^n \mid z_1 = \pm\delta \text{ and } |z'| \leq r\}$ to the boundary $\Gamma_y := \{y \in \mathbb{R}^n \mid y_1 = \pm\delta(1 + |y'|^2) \text{ and } |y'| \leq r\}$, and

$$\frac{\partial \Phi}{\partial z_1}(z) = -\nu(\Phi(z)) \quad \text{on } \Gamma_z, \quad (3.46)$$

$$\left| \frac{\partial y_i}{\partial z_j} \right| \leq C(\bar{r}), \quad \left| \frac{\partial z_i}{\partial y_j} \right| \leq C(\bar{r}), \quad (3.47)$$

where ν is the outward unit normal of Γ_y and C is a positive constant independent of δ .

Proof: Without loss of generality it is enough to show that there exists a transformation $y = \Phi(z)$ which maps $\mathcal{O}_z^+(\delta, r) \doteq \{z \in \mathbb{R}^n \mid 0 \leq z_1 \leq \delta \text{ and } |z'| \leq r\}$ to $\mathcal{O}_y^+(\delta, r) \doteq \{y \in \mathbb{R}^n \mid 0 \leq y_1 \leq \delta(1 + |y'|^2) \text{ and } |y'| \leq r\}$. Denote

$$f(y') = \delta(1 + |y'|^2),$$

then

$$\nu(y') = \nu(f(y'), y') = \frac{(-1, \nabla f(y'))}{\sqrt{1 + |\nabla f(y')|^2}} \quad \text{on } \Gamma_y^+.$$

Define

$$F : (z_1, z') \mapsto (f(z'), z') - (z_1 - \delta)\nu(z'), \quad (3.48)$$

thus F satisfies the conditions (3.45) and (3.46). Plug in f and ν , we have

$$\begin{aligned} F(z_1, z') &= (f(z'), z') - (z_1 - \delta) \frac{(-1, \nabla f(z'))}{\sqrt{1 + |\nabla f(z')|^2}} \\ &= (\delta(1 + |z'|^2), z') - (z_1 - \delta) \frac{(-1, 2\delta z')}{\sqrt{1 + 4\delta^2 z'^2}} \\ &= \left(\frac{z_1}{\sqrt{1 + 4\delta^2 z'^2}}, z' \right) + \delta \left(1 - \frac{1}{\sqrt{1 + 4\delta^2 z'^2}} + |z'|^2, \frac{2(z_1 - \delta)}{\sqrt{1 + 4\delta^2 z'^2}} z' \right). \end{aligned}$$

Construct a fixed cut-off function $0 \leq \rho \leq 1 \in C^\infty(\mathbb{R})$ such that $\rho(t) = 1$ if $t > 3/4$, $\rho(t) = 0$ if $t < 1/4$ and $|\rho'(t)| \leq C$ for all $t \in \mathbb{R}$.

Define

$$\begin{aligned} \Phi(z_1, z') &= \left(\frac{z_1}{\sqrt{1 + 4\delta^2 z'^2}}, z' \right) \\ &\quad + \delta \rho\left(\frac{z_1}{\delta}\right) \left(1 - \frac{1}{\sqrt{1 + 4\delta^2 z'^2}} + |z'|^2, \frac{2(z_1 - \delta)}{\sqrt{1 + 4\delta^2 z'^2}} z' \right) \\ &\doteq \Phi_1(z_1, z') + \delta \rho\left(\frac{z_1}{\delta}\right) \Phi_2(z_1, z'). \end{aligned} \quad (3.49)$$

Thus $y = \Phi(z)$ maps $\mathcal{O}_z^+(\delta, r)$ to $\mathcal{O}_y^+(\delta, r)$ and satisfies conditions (3.45) and (3.46). The only thing need to show is (3.47). By simple calculation we have

$$\begin{aligned} \frac{\partial \Phi}{\partial z_1} &= \frac{\partial \Phi_1}{\partial z_1} + \rho'\left(\frac{z_1}{\delta}\right) \Phi_2 + \delta \rho\left(\frac{z_1}{\delta}\right) \frac{\partial \Phi_2}{\partial z_1} = \frac{\partial \Phi_1}{\partial z_1} + O(|z'|) + O(\delta), \\ \frac{\partial \Phi}{\partial z'} &= \frac{\partial \Phi_1}{\partial z'} + \delta \rho\left(\frac{z_1}{\delta}\right) \frac{\partial \Phi_2}{\partial z'} = \frac{\partial \Phi}{\partial z'} + O(\delta), \end{aligned}$$

Therefore,

$$\begin{aligned} \frac{\partial \Phi}{\partial z} &= \frac{\partial \Phi_1}{\partial z} + O(|z'|) + O(\delta) \\ &= I_{n \times n} + O(\delta^2) + O(|z'|) + O(\delta) \\ &= I_{n \times n} + O(|z'|) + O(\delta), \end{aligned} \quad (3.50)$$

where $|O(|z'|)| \leq C|z'|$ and $|O(\delta)| \leq C\delta$, for some positive constant C independent of $|z'|$ and δ .

Choose $\bar{r} > 0$ small such that $I_{n \times n} + O(\bar{r})$ and its inverse matrix are bounded. Then $\forall 0 < r \leq \bar{r}$ and $\forall \delta > 0$ sufficient small, $\partial_z \Phi$ and its inverse matrix are bounded, i.e. condition (3.47) is proved. \square

Proof of Proposition 3.2: Let $x = \sqrt{\varepsilon}y$, and denote

$$v(y) = u(x) = u(\sqrt{\varepsilon}y). \quad (3.51)$$

Then $\nabla v(y) = \sqrt{\varepsilon}\nabla u(x)$, it is equivalent to show that

$$|\nabla v(y)| \leq C,$$

if

$$y \in \mathcal{O}_y \doteq \left\{ y \in \mathbb{R}^n \mid -\frac{\sqrt{\varepsilon}}{2} - \frac{f(\sqrt{\varepsilon}y')}{\sqrt{\varepsilon}} \leq y_1 \leq \frac{\sqrt{\varepsilon}}{2} + \frac{g(\sqrt{\varepsilon}y')}{\sqrt{\varepsilon}} \text{ and } |y'| \leq \frac{1}{2} \right\}.$$

Since f and g are strictly convex, by the assumptions at the beginning of this section and Lemma 3.8, there exists a transformation $y = \Phi(z) : \mathcal{O}_z(\sqrt{\varepsilon}, r) \rightarrow \mathcal{O}_y(\sqrt{\varepsilon}, r)$. Choose $\bar{r} > 0$ fixed such that $\forall 0 < r \leq \bar{r}$ and $\forall \varepsilon > 0$ sufficient small, we have (3.45), (3.46) and (3.47).

Define

$$w(z) = v(y) = v(\Phi(z)), \quad z \in \mathcal{O}_z(\sqrt{\varepsilon}, r) \quad (3.52)$$

we claim:

(i) $|w| \leq 1$, if $z \in \mathcal{O}_z(\sqrt{\varepsilon}, r)$.

In fact, by definition (3.51), (3.52) of v and w , it follows directly.

(ii) $\frac{\partial w}{\partial z_1} = 0$ on $\Gamma_z = \{z \in \mathbb{R}^n \mid |z_1| = \sqrt{\varepsilon}, |z'| \leq r\}$.

By definition and (3.46), (1.4)

$$\frac{\partial w}{\partial z_1} = \nabla_y v \cdot \frac{\partial \Phi}{\partial z_1} = \sqrt{\varepsilon} \nabla_x u \cdot (-\nu) = -\sqrt{\varepsilon} \frac{\partial u}{\partial \nu} = 0.$$

(iii) $w \in H^1(\mathcal{O}_z)$ satisfied some equation

$$\begin{aligned} \partial_{z_j} \left(a^{ij}(z) \partial_{z_i} w \right) &= 0 & \text{in } \mathcal{O}_z, \\ a^{1\alpha}(z) &= a^{\alpha 1}(z) = 0 & \text{on } |z_1| = \delta, \alpha = 2, 3, \dots, n, \end{aligned}$$

where the $n \times n$ matrix $A(z) := (a^{ij}(z))$ in \mathcal{O}_z is symmetric, $a^{ij}(z) \in C^1(\overline{\mathcal{O}_z})$ and \exists constants $\Lambda \geq \lambda > 0$ such that, for all $\xi \in \mathbb{R}^n$,

$$\lambda |\xi|^2 \leq a^{ij}(z) \xi_i \xi_j \leq \Lambda |\xi|^2 \quad (\forall z \in \mathcal{O}_z).$$

In fact, since u is harmonic, so is v , i.e. v is the minimizer of the energy functional

$$I[\tilde{v}] = \int |\nabla_y \tilde{v}|^2 dy.$$

By definition,

$$\nabla_y v = \nabla_z w \cdot \frac{\partial z}{\partial y},$$

then w is the minimizer of the new energy functional

$$\begin{aligned} I[\tilde{w}] &= \int (\nabla_z \tilde{w} \cdot \frac{\partial z}{\partial y}) (\nabla_z \tilde{w} \cdot \frac{\partial z}{\partial y})^t / |J(z)| \, dz \\ &= \int \nabla_z \tilde{w} \left[\frac{(\partial_y z)(\partial_y z)^t}{|J(z)|} \right] (\nabla_z \tilde{w})^t \, dz. \end{aligned}$$

Let

$$A(z) = (a^{ij}(z)) = \frac{(\partial_z y)^{-1} [(\partial_z y)^{-1}]^t}{|J(z)|} = \frac{[(\partial_z y)^t (\partial_z y)]^{-1}}{|J(z)|}.$$

By (3.46) we have for $\alpha = 2, 3, \dots, n$,

$$\partial_{z_1} y \cdot \partial_{z_\alpha} y = 0, \quad \text{on } |z_1| = \delta.$$

Therefore, for $\alpha = 2, 3, \dots, n$

$$a^{1\alpha}(z) = a^{\alpha 1}(z) = 0 \quad \text{on } |z_1| = \delta.$$

Hence, by (3.47), we derive claim (iii).

Thus, w satisfies all conditions in Lemma 3.7. Therefore we have

$$|\nabla_z w| \leq C \quad \text{in } \mathcal{O}_z(\sqrt{\varepsilon}, \frac{r}{2}).$$

Hence, by (3.47), in $\mathcal{O}_y(\sqrt{\varepsilon}, r/2)$

$$|\nabla_y v| = |\nabla_z w(\partial_y z)| \leq C.$$

The proof is complete. □

Proof of Theorem 1.6: We divided $\tilde{\Omega}$ into two pieces $B(0, \frac{\sqrt{\varepsilon}}{2}) \cap \tilde{\Omega}$ and $\tilde{\Omega}/B(0, \frac{\sqrt{\varepsilon}}{2})$ and estimate $|\nabla u|$ separately.

In $B(0, \frac{\sqrt{\varepsilon}}{2}) \cap \tilde{\Omega}$, by Proposition 3.2, we have $|\nabla u| \leq \frac{C}{\sqrt{\varepsilon}}$.

In $\tilde{\Omega}/B(0, \frac{\sqrt{\varepsilon}}{2})$, for any point $x = (x_1, x')$ in it, instead of rescaling $x = \sqrt{\varepsilon}y$ in the first step in the proof of Proposition 3.2, let

$$x = \sqrt{|x'|^2 + \varepsilon} \cdot y,$$

and follow the same proof the Proposition 3.2, we have

$$|\nabla u| \leq \frac{C}{\sqrt{|x'|^2 + \varepsilon}} \leq \frac{C}{\sqrt{\varepsilon}}.$$

We complete the proof of Theorem 1.6. \square

Remark 3.2 *By the proof of Theorem 1.6, indeed, we have derived a more precise point wise gradient estimate*

$$|\nabla u(x_1, x')| \leq \frac{C}{\sqrt{|x'|^2 + \varepsilon}}.$$

3.3 Appendix B: Some elementary results for the insulated conductivity problem

Assume that in \mathbb{R}^n , Ω and ω are bounded open sets with $C^{2,\alpha}$ boundaries, $0 < \alpha < 1$, satisfying

$$\bar{\omega} = \bigcup_{s=1}^m \bar{\omega}_s \subset \Omega,$$

where $\{\omega_s\}$ are connected components of ω . Clearly, $m < \infty$ and ω_s is open for all $1 \leq s \leq m$. Given $\varphi \in C^2(\partial\Omega)$, the conductivity problem we consider is the following transmission problem with Dirichlet boundary condition:

$$\begin{cases} \partial_{x_j} \left\{ \left[(ka_1^{ij}(x) - a_2^{ij}(x))\chi_\omega + a_2^{ij}(x) \right] \partial_{x_i} u_k \right\} = 0 & \text{in } \Omega, \\ u_k = \varphi & \text{on } \partial\Omega, \end{cases} \quad (3.53)$$

where $k = 1, \frac{1}{2}, \frac{1}{3}, \dots$, and χ_ω is the characteristic function of ω .

The $n \times n$ matrixes $A_1(x) := (a_1^{ij}(x))$ in ω , $A_2(x) := (a_2^{ij}(x))$ in $\Omega \setminus \overline{\omega}$ are symmetric and \exists a constant $\Lambda \geq \lambda > 0$ such that

$$\lambda |\xi|^2 \leq a_1^{ij}(x) \xi_i \xi_j \leq \Lambda |\xi|^2 \quad (\forall x \in \omega), \quad \lambda |\xi|^2 \leq a_2^{ij}(x) \xi_i \xi_j \leq \Lambda |\xi|^2 \quad (\forall x \in \Omega \setminus \omega)$$

for all $\xi \in \mathbb{R}^n$ and $a_1^{ij}(x) \in C^2(\overline{\omega})$, $a_2^{ij}(x) \in C^2(\overline{\Omega \setminus \omega})$.

Equation (3.53) can be rewritten in the following form to emphasize the transmission condition on $\partial\omega$:

$$\begin{cases} \partial_{x_j} (a_1^{ij}(x) \partial_{x_i} u_k) = 0 & \text{in } \omega, \\ \partial_{x_j} (a_2^{ij}(x) \partial_{x_i} u_k) = 0 & \text{in } \Omega \setminus \overline{\omega}, \\ u_k|_+ = u_k|_-, & \text{on } \partial\omega, \\ a_2^{ij}(x) \partial_{x_i} u_k \nu_j|_+ = k a_1^{ij}(x) \partial_{x_i} u_k \nu_j|_- & \text{on } \partial\omega, \\ u_k = \varphi & \text{on } \partial\Omega. \end{cases} \quad (3.54)$$

We list the following results which are well known and omit the proofs.

Theorem 3.2 *If $u_k \in H^1(\Omega)$ is a solution of equation (3.53), then $u_k \in C^1(\overline{\Omega \setminus \omega}) \cap C^1(\overline{\omega})$ and satisfies equation (3.54).*

If $u_k \in C^1(\overline{\Omega \setminus \omega}) \cap C^1(\overline{\omega})$ is a solution of equation (3.54), then $u_k \in H^1(\Omega)$ and satisfies equation (3.53).

Theorem 3.3 *There exists at most one solution $u_k \in H^1(\Omega)$ to equation (3.53).*

The existence of the solution can be obtained by using the variational method. For every k , we define the energy functional

$$I_k[v] := \frac{k}{2} \int_{\omega} a_1^{ij}(x) \partial_{x_i} v \partial_{x_j} v + \frac{1}{2} \int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} v \partial_{x_j} v, \quad (3.55)$$

where v belongs to the set

$$H_{\varphi}^1(\Omega) := \{v \in H^1(\Omega) \mid v = \varphi \text{ on } \partial\Omega\}.$$

Theorem 3.4 *For every k , there exists a minimizer $u_k \in H^1(\Omega)$ satisfying*

$$I_k[u_k] = \min_{v \in H_{\varphi}^1(\Omega)} I_k[v].$$

Moreover, $u_k \in H^1(\Omega)$ is a solution of equation (3.53).

Comparing equation (3.54), when $k = 0$, the insulated conducting problem turns out to be:

$$\begin{cases} \partial_{x_j} \left(a_2^{ij}(x) \partial_{x_i} u_0 \right) = 0 & \text{in } \Omega \setminus \overline{\omega}, \\ a_2^{ij}(x) \partial_{x_i} u_0 \nu_j|_+ = 0 & \text{on } \partial\omega, \\ u_0 = \varphi & \text{on } \partial\Omega, \\ \partial_{x_j} \left(a_1^{ij}(x) \partial_{x_i} u_0 \right) = 0 & \text{in } \omega, \\ u_0|_+ = u_0|_-, & \text{on } \partial\omega. \end{cases} \quad (3.56)$$

We also have similar results:

Theorem 3.5 *If $u_0 \in H^1(\Omega)$ satisfies equation (3.56), then $u_0 \in C^1(\overline{\Omega \setminus \omega}) \cap C^1(\overline{\omega})$.*

Theorem 3.6 *There exists at most one solution $u_0 \in H^1(\Omega) \cap C^1(\overline{\Omega \setminus \omega}) \cap C^1(\overline{\omega})$ of equation (3.56).*

Proof: It is equivalent to showing that if $\varphi = 0$, equation (3.56) only has the solution $u_0 \equiv 0$. Integrating by parts in the first line of equation (3.56), we have

$$\begin{aligned} 0 &= - \int_{\Omega \setminus \overline{\omega}} \partial_{x_j} \left(a_2^{ij}(x) \partial_{x_i} u_0 \right) \cdot u_0 \\ &= \int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} u_0 \partial_{x_j} u_0 - \int_{\partial\Omega} u_0 \cdot a_2^{ij}(x) \partial_{x_i} u_0 \nu_j|_- + \int_{\partial\omega} u_0 \cdot a_2^{ij}(x) \partial_{x_i} u_0 \nu_j|_+ \\ &\geq \lambda \int_{\Omega \setminus \overline{\omega}} |\nabla u_0|^2 - \int_{\partial\Omega} \varphi \cdot a_2^{ij}(x) \partial_{x_i} u_0 \nu_j|_- \\ &= \lambda \int_{\Omega \setminus \overline{\omega}} |\nabla u_0|^2. \end{aligned}$$

Thus $\nabla u_0 = 0$ in $\Omega \setminus \overline{\omega}$. And since $u_0 = \varphi = 0$ on $\partial\Omega$, we have $u_0 \equiv 0$ in $\Omega \setminus \overline{\omega}$. Since $u_0|_+ = u_0|_-$ on $\partial\omega$ and $\partial_{x_j} \left(a_1^{ij}(x) \partial_{x_i} u_0 \right) = 0$ on $\overline{\omega}$, we get $u_0 = 0$ on $\overline{\omega}$. Hence $u_0 \equiv 0$ in Ω , i.e. $u_0 \equiv 0$ is the only solution of (3.56) when $\varphi = 0$. \square

Define the energy functional

$$I_0[v] := \frac{1}{2} \int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} v \partial_{x_j} v, \quad (3.57)$$

where v belongs to the set

$$\mathcal{A}_0 := \{v \in H_\varphi^1(\Omega) \mid \partial_{x_j} \left(a_1^{ij}(x) \partial_{x_i} v \right) = 0 \text{ in } \omega\}.$$

Theorem 3.7 *There exists a minimizer $u_0 \in \mathcal{A}_0$ satisfying*

$$I_0[u_0] = \min_{v \in \mathcal{A}_0} I_0[v].$$

Moreover, $u_0 \in H^1(\Omega) \cap C^1(\overline{\Omega \setminus \omega}) \cap C^1(\overline{\omega})$ is a solution of equation (3.56).

Proof: By the lower-semi continuity of I_0 and the weakly closed property of \mathcal{A}_0 , it is easy to see that the minimizer $u_0 \in \mathcal{A}_0$ exists and satisfies $\partial_{x_j} \left(a_2^{ij}(x) \partial_{x_i} u_0 \right) = 0$ in $\Omega \setminus \overline{\omega}$. The only thing which needs to be shown is the second line in equation (3.56), i.e.

$$a_2^{ij}(x) \partial_{x_i} u_0 \nu_j|_+ = 0 \text{ on } \partial\omega.$$

In fact, since u_0 is a minimizer, for any $\phi \in C_c^\infty(\Omega)$ satisfying $\partial_{x_j} \left(a_1^{ij}(x) \partial_{x_i} u_0 \right) = 0$ on $\overline{\omega}_s$ and $\phi \equiv 0$ on $\overline{\omega}_t$ ($t \neq s$), let

$$i(t) := I_0[u_0 + t\phi] \quad (t \in \mathbb{R}),$$

we have

$$i'(0) := \frac{di}{dt} \Big|_{t=0} = \int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} u_0 \phi_{x_j} = 0.$$

Therefore

$$\begin{aligned} 0 &= - \int_{\Omega \setminus \overline{\omega}} \partial_{x_j} \left(a_2^{ij}(x) \partial_{x_i} u_0 \right) \phi = \int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} u_0 \phi_{x_j} + \int_{\partial\omega} \phi \cdot a_2^{ij}(x) \partial_{x_i} u_0 \nu_j|_+ \\ &= \int_{\partial\omega_s} \phi \cdot a_2^{ij}(x) \partial_{x_i} u_0 \nu_j|_+, \end{aligned}$$

for any $\phi \in C_c^\infty(\Omega)$. Hence, $a_2^{ij}(x) \partial_{x_i} u_0 \nu_j|_+ = 0$ on $\partial\omega_s$ for any s . \square

Finally, we give the relationship between u_k and u_0 .

Theorem 3.8 *Let u_k and u_0 in $H^1(\Omega)$ be the solutions of equations (3.54) and (3.56), respectively. Then*

$$u_k \rightharpoonup u_0 \text{ in } H^1(\Omega), \quad \text{as } k \rightarrow 0,$$

and

$$\lim_{k \rightarrow 0} I_k[u_k] = I_0[u_0],$$

where I_k and I_0 are defined as (3.55) and (3.57).

Proof: Step 1. By the uniqueness of the solution to equation (3.56), we only need to show that there exists a weak limit u_0 of a subsequence of $\{u_k\}$ in $H^1(\Omega)$ and u_0 is the solution of equation (3.56).

(1) To show that after passing to a subsequence, u_k weakly converges in $H^1(\Omega)$ to some u_0^* .

Since u_k is the minimizer of I_k in $H_\varphi^1(\Omega)$ and u_0 is the minimizer of I_0 in \mathcal{A}_0 , we have

$$\begin{aligned} \frac{\lambda k}{2} \|\nabla u_k\|_{L^2(\omega)} + I_0[u_0] &\leq \frac{k}{2} \int_{\omega} a_1^{ij}(x) \partial_{x_i} u_k \partial_{x_j} u_k + I_0[u_0] \\ &\leq \frac{k}{2} \int_{\omega} a_1^{ij}(x) \partial_{x_i} u_k \partial_{x_j} u_k + I_0[u_k] = I_k[u_k] \\ &\leq I_k[u_0] = \frac{k}{2} \int_{\omega} a_1^{ij}(x) \partial_{x_i} u_0 \partial_{x_j} u_0 + I_0[u_0], \\ &\leq \frac{\Lambda k}{2} \|\nabla u_0\|_{L^2(\omega)} + I_0[u_0], \end{aligned}$$

i.e.

$$\|\nabla u_k\|_{L^2(\omega)} \leq \frac{\Lambda}{\lambda} \|\nabla u_0\|_{L^2(\omega)} \leq \frac{\Lambda}{\lambda} \|\nabla u_0\|_{L^2(\Omega)}.$$

On the other hand,

$$\frac{\lambda}{2} \|\nabla u_k\|_{L^2(\Omega \setminus \overline{\omega})} \leq I_k[u_k] \leq I_k[u] \leq \frac{\Lambda}{2} \|\nabla u_0\|_{L^2(\Omega)},$$

i.e.

$$\|\nabla u_k\|_{L^2(\Omega \setminus \overline{\omega})} \leq \frac{\Lambda}{\lambda} \|\nabla u_0\|_{L^2(\Omega)}.$$

Therefore,

$$\|\nabla u_k\|_{L^2(\Omega)} = \|\nabla u_k\|_{L^2(\omega)} + \|\nabla u_k\|_{L^2(\Omega \setminus \overline{\omega})} \leq \frac{2\Lambda}{\lambda} \|\nabla u_0\|_{L^2(\Omega)} \doteq \overline{M},$$

where \overline{M} is independent of k .

Since $u_k = \varphi$ on $\partial\Omega$ and $\sup_k \|u_k\|_{H^1(\Omega)} < \infty$, we have $u_k \rightharpoonup u_0^*$ in $H_\varphi^1(\Omega)$ as $k \rightarrow \infty$.

(2) To show that u_0^* is also a solution of equation (3.56). Therefore, $u_0^* \equiv u_0$.

In fact, we only need to prove the following three conditions:

$$\partial_{x_j} \left(a_2^{ij}(x) \partial_{x_i} u_0^* \right) = 0 \quad \text{in } \Omega \setminus \overline{\omega}, \quad (3.58)$$

$$\partial_{x_j} \left(a_1^{ij}(x) \partial_{x_i} u_0^* \right) = 0 \quad \text{in } \omega, \quad (3.59)$$

$$a_2^{ij}(x) \partial_{x_i} u_0^* \nu_j|_+ = 0 \quad \text{on } \partial\omega. \quad (3.60)$$

(i) For every k , since $u_k \in H^1(\Omega)$ is the solution of equation (3.53), then

$\forall \phi \in C_c^\infty(\Omega)$, we have

$$k \int_{\omega} a_1^{ij}(x) \partial_{x_i} u_k \phi_{x_j} + \int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} u_k \phi_{x_j} = 0.$$

Thus, $\forall \phi \in C_c^\infty(\Omega \setminus \overline{\omega}) \subset C_c^\infty(\Omega)$,

$$0 = \int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} u_k \phi_{x_j} \longrightarrow \int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} u_0^* \phi_{x_j},$$

since $u_k \rightharpoonup u_0^*$ in $H_\varphi^1(\Omega) \subset H^1(\Omega)$.

Therefore,

$$\int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} u_0^* \phi_{x_j} = 0, \quad \forall \phi \in C_c^\infty(\Omega \setminus \overline{\omega}),$$

i.e. (3.58).

(ii) Similar to (i), $\forall \phi \in C_c^\infty(\omega) \subset C_c^\infty(\Omega)$, we obtain (3.59).

(iii) By (i) and (ii), $u_0^* \in C^2(\overline{\Omega \setminus \omega})$. For any $\varrho \in C_c^2(\overline{\Omega})$, by Green's Identity, we have the following:

$$\begin{aligned} 0 &= - \int_{\Omega \setminus \overline{\omega}} \partial_{x_j} \left(a_2^{ij}(x) \partial_{x_i} u_k \right) \varrho \\ &= \int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} u_k \partial_{x_j} \varrho - \int_{\partial \Omega} \varrho \cdot a_2^{ij}(x) \partial_{x_i} u_k \nu_j|_- + \int_{\partial \omega} \varrho \cdot a_2^{ij}(x) \partial_{x_i} u_k \nu_j|_+ \\ &= \int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} u_k \partial_{x_j} \varrho + k \int_{\partial \omega} \varrho \cdot a_1^{ij}(x) \partial_{x_i} u_k \nu_j|_- \\ &\longrightarrow \int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} u_0^* \partial_{x_j} \varrho, \end{aligned}$$

Since $u_k \rightharpoonup u_0^*$ in $H^1(\Omega)$.

Similarly,

$$0 = - \int_{\Omega \setminus \overline{\omega}} \partial_{x_j} \left(a_2^{ij}(x) \partial_{x_i} u_0^* \right) \varrho = \int_{\Omega \setminus \overline{\omega}} a_2^{ij}(x) \partial_{x_i} u_0^* \partial_{x_j} \varrho + \int_{\partial \omega} \varrho \cdot a_2^{ij}(x) \partial_{x_i} u_0^* \nu_j|_+.$$

Thus,

$$\int_{\partial \omega} \varrho \cdot a_2^{ij}(x) \partial_{x_i} u_0^* \nu_j|_+ = 0,$$

for any $\varrho \in C_c^2(\overline{\Omega})$. Therefore, we obtain (3.60) and finish the proof of the first part.

Step 2. Since u_k is the minimizer of I_k in $H^1_\varphi(\Omega)$ and u_0 is the minimizer of I_0 in \mathcal{A}_0 , we have

$$\begin{aligned} I_0[u_0] &\leq I_0[u_k] \leq \frac{k}{2} \int_{\omega} a_1^{ij}(x) \partial_{x_i} u_k \partial_{x_j} u_k + I_0[u_k] = I_k[u_k] \\ &\leq I_k[u_0] = \frac{k}{2} \int_{\omega} a_1^{ij}(x) \partial_{x_i} u_0 \partial_{x_j} u_0 + I_0[u_0] \longrightarrow I_0[u_0], \end{aligned}$$

Therefore,

$$\lim_{k \rightarrow 0} I_k[u_k] = I_0[u_0].$$

□

References

- [1] H. Ammari, M. Asch, H. Kang, *Boundary voltage perturbations caused by small conductivity inhomogeneities nearly touching the boundary*, Adv. Appl. Math. 35 (2005), 368-391.
- [2] H. Ammari, G. Dassios, H. Kang and M. Lim, *Estimates for the electric field in the presence of adjacent perfectly conducting spheres*, Quat. Appl. Math. 65 (2007), 339-355.
- [3] H. Ammari and H. Kang, *Reconstruction of small inhomogeneities from boundary measurements*, Lecture Notes in Mathematics, vol. 1846, Springer-Verlag, 2004.
- [4] H. Ammari, H. Kang, *Polarization and Moment Tensors: with Applications to Inverse Problems and Effective Medium Theory*, Applied Mathematical Sciences, vol. 162, Springer-Verlag, New York, 2007.
- [5] H. Ammari, H. Kang, E. Kim, M. Lim, *Reconstruction of closely spaced small inclusions*, SIAM J. Numer. Anal. 42 (2005), 2408-2428.
- [6] H. Ammari, H. Kang, H. Lee, J. Lee and M. Lim, *Optimal estimates for the Electrical Field in Two Dimensions*, J. Math. Pures Appl. 88 (2007), 307-324.
- [7] H. Ammari, H. Kang and M. Lim, *Gradient estimates for solutions to the conductivity problem*, Math. Ann. 332 (2005), 277-286.
- [8] H. Ammari, H. Kang, K. Touibi, *Approximation of a conductivity inclusion close to a planar surface*, J. Appl. Math. Phys. 57 (2006) 234-243.
- [9] I. Babuska, B. Anderson, P.J. Smith, and K. Levin, *Damage analysis of fiber composites. I. Statistical analysis on fiber scale*, Comput. Methods Appl. Mech. Engrg. 172 (1999), 27-77.
- [10] M. F. Ben Hensen and E. Bonnetier, *Asymptotic formulas for the voltage potential in a composite medium containing closer or touching disks of small diameter*, Multiscale Model. Simul., 4 (2005), 250-277.
- [11] E.S. Bao, Y.Y. Li and B. Yin, *Gradient estimates for the perfect conductivity problem*, (arXiv:math.AP/0612520), Arch. Rational Mech. and Anal., to appear.
- [12] E.S. Bao, Y.Y. Li and B. Yin, *Gradient estimates for the perfect and insulated conductivity problem and elliptic systems*, in preparation.
- [13] E.S. Bao and B. Yin, *Gradient estimates for the conductivity problem and elliptic systems with inclusions close to boundary*, in preparation.
- [14] E. Bonnetier and M. Vogelius, *An elliptic regularity result for a composite medium with "touching" fibers of circular cross-section*, SIAM J. Math. Anal. 31 (2000), 651-677.

- [15] B. Budiansky and G.F. Carrier, *High shear stresses in stiff fiber composites*, J. App. Mech. 51 (1984), 733-735.
- [16] A. Charalambopoulos, G. Dassios, and M. Hadjinicolaou, *An analytic solution for lowfrequency scattering by two soft spheres*, SIAM J. Appl. Math., 58 (1998), 370-386.
- [17] G.B. Folland, *Introduction to Partial Differential Equations*, Princeton University Press, Princeton, NJ, 1976.
- [18] D. Gilbarg and N. S. Trudinger, *Elliptic partial differential equations of second order*, Reprint of the 1998 edition, Classics in Mathematics, Springer-Verlag, Berlin, 2001.
- [19] J. G. Goree and H. B. Wilson Jr., *Transverse shear loading in an elastic matrix containing two circular cylindrical inclusions*, Trans. ASEM J. Appl. Mech., 34 (1967), 511-513.
- [20] J. D. Jackson, *Classical Electrodynamics*, 3rd ed., Wiley, New York, 1999.
- [21] H. Kang, J.K. Seo, *Layer potential technique for the inverse conductivity problem*, Inverse Problems 12 (1996), 267-278.
- [22] H. Kang, J.K. Seo, *Recent progress in the inverse conductivity problem with single measurement*, in: *Inverse Problems and Related Fields*, CRC Press, Boca Raton, FL, 2000, 69-80.
- [23] J.B. Keller, *Conductivity of a medium containing a dense array of perfectly conducting spheres or cylinders or nonconducting cylinders*, J. Appl. Phys., 3 (1963), 991-993.
- [24] J.B. Keller, *Stresses in narrow regions*, Trans. ASME J. Appl. Mech., 60 (1993), 1054-1056.
- [25] Y.Y. Li and L. Nirenberg, *Estimates for elliptic system from composite material*, Comm. Pure Appl. Math. 56 (2003), 892-925.
- [26] Y.Y. Li and M. Vogelius, *Gradient estimates for solution to divergence form elliptic equation with discontinuous coefficients*, Arch. Rational Mech. Anal. 153 (2000), 91-151.
- [27] X. Markenscoff, *Stress amplification in vanishingly small geometries*, Computational Mechanics 19 (1996), 77-83.
- [28] P. Moon and D.E. Spencer, *Field Theory Handbook*, 2nd Ed. Springer-Verlag, Berlin, 1988.
- [29] G.C. Verchota, *Layer potentials and boundary value problems for Laplaces equation in Lipschitz domains*, J. Funct. Anal. 59 (1984), 572-611.
- [30] M. Vogelius and D. Volkov, *Asymptotic formulas for perturbations in the electromagnetic fields due to the presence of inhomogeneities*, M2AN Math. Model. Numer. Anal., 34 (2000), 723-748.
- [31] K. Yun, *Estimates for electric fields blown up between closely adjacent conductors with arbitrary shape*, SIAM J. on Applied Math. 67 (2007), 714-730.

- [32] K. Yun, *Optimal bound on high stresses occurring between stiff fibers with arbitrary shaped cross-sections*, preprint.

Vita

ShiTing Bao

- 2008** Ph. D. in Mathematics, Rutgers University
- 1995-2002** B. Sc. in Mathematics, Special Class for the Gifted Young (SCGY), University of Science and Technology of China (USTC)
- 1995** Graduated from Suzhou Middle School in Jiangsu Province of China
- 2002-2008** Teaching assistant, Department of Mathematics, Rutgers University